

Method for the specific rapid detection of beverage-spoiling microorganisms

5 The invention is related to a method for the specific fast detection of drink-spoiling microorganisms by *in situ*-hybridization. Moreover, the invention is related to specific oligonucleotide probes which are used in the course of the method for detection as well as kits which contain these oligonucleotide probes.

10 Under the generic clause "non-alcoholic drinks" groups of beverages are summarized like fruit juices, fruit nectars, fruit concentrates, mashed fruits, soft drinks and waters.

Basically non-alcoholic drinks can, due to their diverse/varying composition of nutrients and growth stimulating substances, be classified as potentially endangered  
15 by the growth of a large variety of microorganisms.

According to present knowledge mainly yeasts, molds, lactic acid bacteria, acetic acid bacteria, bacilli and alicyclobacilli are found in non-alcoholic drinks and are thus described as "drink-spoiling" microorganisms.  
20

In general contaminations with these microorganisms do not lead to health defects of the consumer but are associated with turbidity, changes of taste and smell within the endproduct and cause high economic losses for the producing industry by image damage based thereon.  
25

Based on the naturally high concentrations of fruit acids and a corresponding low pH-value (a pH range from 2.5 to 4.5) in fruit juices and fruit nectars only acidophilic or acidotolerant microorganisms (such as lactic acid bacteria, alicyclobacilli, acid tolerant yeast and mold species) can grow and subsequently lead  
30 to a deterioration of these beverages.

A measure for restricting spoilage due to microorganisms is carbonisation of beverages. This method is commonly used for the production of soft drinks. By the

addition of CO<sub>2</sub> almost anaerobic conditions are created in the product and only micro-aerophilic, facultatively anaerobic and anaerobic microorganisms (such as lactic acid bacteria, acetic acid bacteria and yeasts) are able to tolerate this environment.

5

Non-carbonated beverages are in most cases pasteurised in order to assure a long stability and quality of these products. By pasteurisation all vegetative microorganisms should be killed in a manner as comprehensive as possible. However, spores formed by bacilli or alicyclobacilli are not eliminated by this measure. Furthermore, some mold species are able to sustain this process without damage and subsequently create product damages.

15 A crucial factor for guaranteeing the biological quality of the beverages is the search for the cause of contamination in order to finally eliminate the same. In general, two ways of contamination are distinguished: contaminations are characterised as primary contaminations when microorganisms are introduced into the process by the raw material or by contamination within the process.

20 Secondary contaminations are those which appear in the filling area after the actual production of the beverage.

The challenge which arises by these different factors for the microbiological quality control, resides in the comprehensive and fast identification of all cells present in the product in order to be able to initiate corresponding counter measures as fast as possible.

25

Until now conventional detection of drink-spoiling microorganisms is performed by a several days lasting enrichment of the sample in a selective culture medium followed by light microscopy. Furthermore, for the accurate identification of the

drink-spoiling microorganism further physiological tests (like Gram-staining, sugar consumption tests) need to be carried out.

5 The disadvantages of this solely cultivation-based method are the long duration of the analysis, which cause significant logistic costs in beverage-producing companies. Furthermore, there is the threat of significant image loss for said company, if, after the delivery of products whose microbiological findings had not yet been unequivocally stated, contaminationen are realised and draw-back actions of the spoiled product batches are required.

10

In the following the drink-spoiling microorganisms and their state of the art detection is described in detail.

#### Yeasts and molds:

15

Microorganisms which can survive heat treatment and cause subsequently problems in the beverages are mainly the molds *Byssochlamys fulva* and *B. nivea*, *Neosartorya fischeri* and *Talaromyces flavus* as well as some yeasts. In carbonated drinks mainly the acid-tolerant, fermentative members of yeasts (*Saccharomyces spp.*, *Dekkera spp.* and *Zygosaccharomyces bailii*) are dominating. Besides the threat of product damage based on taste alterations and turbidity caused by these “fermentative yeasts” there is a potential danger of occasional bursts of the filled bottles.

20

The detection of yeasts and molds is currently performed by cultivation on  
25 corresponding culture media (e.g. SSL-bouillon, OFS-medium, malt-dextrose-medium, wort-agar) and needs between 2 and 7 days. A detection on genus or even species level is very time-consuming and is normally not performed.

#### Lactic acid bacteria

The members of lactic acid bacteria are Gram-positive, non spore-forming, catalase-negative rods and cocci which are characterised by a very high nutrient demand (above all vitamins, amino acids, purines and pyrimidines). As indicated by the name all lactic acid bacteria are able to produce lactic acid as fermentation product.

5

Due to their anaerobic growth and for anaerobic microorganisms atypical high tolerance and insensitivity against oxygen they are described as aerotolerant anaerobics.

- 10 Up until now the genera *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Oenococcus*, *Carnobacterium*, *Bifidobacterium*, *Enterococcus*, *Pediococcus*, *Weissella* and *Streptococcus* are referred to as "lactic acid bacteria".

- 15 Lactic acid bacteria play an ambivalent role in the food industry. On the one side their presence is wished and indispensable in some processes such as, e.g., the production of sauerkraut. On the other side their presence in beer or fruit juices can lead to a deterioration of the products. The growth of these bacteria is manifested mainly by turbidity, acidification and formation of gas and slime.

- 20 In the non-alcoholic drinks industry mainly the bacterial genera *Leuconostoc*, *Lactococcus*, *Lactobacillus*, *Oenococcus*, *Weissella* and *Pediococcus* are relevant as contaminants. Lactic acid bacteria are detected by a 5 to 7 days incubation at 25°C on MRS agar (pH 5.7).

- 25 Acetic acid bacteria

Bacteria of the genera *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter* and *Acidomonas* are described with the trivial name "acetic acid bacteria". Bacteria of these genera are gram-negative, obligate aerobic, oxidase-negative rods whose optimum growth temperature is at 30°C. Acetic acid bacteria are able to grow also at

pH values of 2.2 to 3.0 and, therefore, can produce product damages in beverages having this pH value.

Phylogenetically, bacteria of this genus are members of the *Alphaproteobacteria*.

5

The product damages mainly goes along with turbidity and alteration of the taste by the formation of acetic acid and gluconic acid. For the detection of acetic acid bacteria mainly ACM-agar (incubation time: 14 days) and DSM-agar (incubation time: 3 to 5 days) have proved themselves.

10

#### Bacilli:

Bacilli are Gram-positive aerobic, partly facultatively anaerobic, mostly catalase-positive spore-forming rods. Up until now *Bacillus coagulans* was mainly identified as spoilage microorganism in the non-alcoholic beverage industry.

15

The detection is performed by plating the sample on dextrose-caseine-peptone agar or yeast extract-peptone dextrose starch agar and subsequent incubation at 55°C (incubation time: 3 days). In order to activate the spores and to achieve a germination of the spores of *B. coagulans*, respectively, a heat treatment of the sample is recommended at 80°C for 10 min. before the actual incubation.

20

#### Alicyclobacilli:

Alicyclobacilli are Gram-positive, aerobic, thermophilic and catalase-positive spore-forming rods. Members of this genus produce  $\gamma$ -alicyclic fatty acids as main fatty acids. Up until now *Alicyclobacillus acidoterrestris* was mainly identified in the non-alcoholic beverage industry as spoilage organism. In rare cases also *A. acidocaldarius* and *A. acidiphilus* were identified in spoiled beverages.

25

The optimum range of the growth temperature for *Alicyclobacillus* spp. is between 26 and 55°C. The pH range where bacteria of this genus can grow, is between 2.2 and 5.8.

- 5 The growth of *A. acidoterrestris* leads to spoilage in fruit juices, which is manifested as alteration of the smell and taste due to the formation of guaiacol and di-bromophenol. A contamination with this organism proceeds mostly in a non-apparent way, which means that only in rare cases a turbidity is seen in infected beverages.
- 10 Alicyclobacilli can be detected by cultivation for several days at 44 – 46°C on orange serum agar, potato dextrose agar, K-agar, YSG-agar or BAM-agar. Furthermore, for the exact confirmation of the finding a set of physiological tests is necessary. In order to activate the spores and to achieve a germination of the spores of *Alicyclobacillus* ssp., respectively, heat treatment of the sample is recommended at 80°C for 10 min.
- 15 before the actual incubation.

The routine detection methods for drink-spoiling microorganisms used so far, are very protracted and are partly too inaccurate and thus prevent fast and effective counter measures in order to save the contaminated product. The inaccuracy of the

20 detection arises from a missing differentiation up to genus and/or species level.

As a logical consequence of the difficulties presented by traditional cultivation methods for the detection of drink-spoiling microorganisms, detection methods on the basis of nucleic acids are suitable for the fast, safe and specific identification of

25 spoilage microorganisms in non-alcoholic beverages.

In PCR, which is polymerase chain reaction, a characteristic piece of the respective bacterial genome is amplified with specific primers. If the primer finds its target site, a million-fold amplification of a piece of the inherited material occurs. In the

30 following analysis, for example by an agarose gel separating DNA fragments, a

qualitative evaluation can take place. In the most simple case this leads to the conclusion that target sites for the primers used were present in the tested sample. Further conclusions are not possible; these target sites can originate from both a living bacterium and a dead bacterium, or from naked DNA. Since the PCR reaction is positive also in the presence of a dead bacterium or naked DNA, this often leads to false-positive results. A further refinement of this technique is the quantitative PCR which aims at establishing a correlation between the amount of bacteria present and the amount of amplified DNA. Advantages of the PCR are its high specificity, its ease of application and its low expenditure of time. Its main disadvantages are its high susceptibility to contamination and therefore false-positive results, as well as the aforementioned lacking possibility to discriminate between viable and dead cells, and naked DNA, respectively.

A unique approach to combine the specificity of molecularbiological methods such as PCR and the possibility of visualizing bacteria, which is provided by the antibody methods, is the method of fluorescence *in situ* hybridization (FISH; R.I. Amann, W. Ludwig and K.-H. Schleifer, 1995. Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. Microbiol. Rev. 59, p. 143-169). Using this method bacteria species, genera or groups can be identified and visualized with high specificity.

The FISH technique is based on the fact that in cells of microorganism there are certain molecules which have been mutated only to a small extent in the course of evolution because of their essential function. These are the 16S and the 23S ribosomal ribonucleic acid (rRNA). Both are components of the ribosomes, the sites of protein biosynthesis, and can serve as specific markers on account of their ubiquitous distribution, their size and their structural and functional constancy (Woese, C.R., 1987. Bacterial evolution. Microbiol. Rev. 51, p. 221-271). Based on a comparative sequence analysis, phylogenetic relationships can be established based on these data alone. For this purpose, the sequence data have to be brought into an

alignment. In the alignment, which is based on the knowledge about the secondary structure and tertiary structure of these macromolecules, the homologous positions of the ribosomal nucleic acids are brought in line with each other.

- 5 Based on these data, phylogenetic calculations can be made. The use of the most modern computer technology allows to perform even large-scale calculations fast and effectively, as well as to set up large databases which contain the alignment sequences of the 16S, 18S, 23S and 26S rRNA. Due to the fast access to this data material, newly acquired sequences can be phylogenetically analyzed within a short
- 10 time. These rRNA databases can be used to design species-specific and genus-specific gene probes. Hereby all available rRNA sequences are compared with each other and probes are designed for specific sequence sites, which specifically target a specific species, genus or group of bacteria.
- 15 In the FISH (fluorescence *in situ* hybridization) technique, these gene probes which are complementary to a certain region on the ribosomal target sequence, are introduced into the cell. The gene probes are generally small, 16-20 bases long, single-stranded deoxyribonucleic acid pieces and are directed against a target region which is characteristic for a bacterial species or a bacterial group. If a
- 20 fluorescently labeled gene probe finds its target sequence in a cell of a microorganisms, it binds to it and the cells can be detected by means of a fluorescence microscope because of their fluorescence.

The FISH analysis is always performed on a slide, as for the evaluation the bacteria

25 are visualized, i. e. rendered visible, by irradiation with high-energy light. But herein lies one of the disadvantages of the classical FISH analysis: because by definition only comparatively small volumina can be analyzed on a slide, the sensitivity of the method is not satisfying and not sufficient for a reliable analysis.



The present invention thus combines the advantages of the classical FISH analysis with those of cultivation. A comparatively short cultivation step ensures that the bacteria to be detected are present in sufficient numbers before the bacteria are detected using specific FISH.

5

The practising of the methods described in the present application for the specific detection of drink-spoiling yeasts of the genera *Zygosaccharomyces*, *Hanseniaspora*, *Candida*, *Brettanomyces*, *Dekkera*, *Pichia*, *Saccharomyces* and *Saccharomycodes* in particular the species *Zygosaccharomyces bailii*, *Z. mellis*, *Z. rouxii*, *Z. bisporus*, *Z. fermentati*, *Z. microellipsoides*, *Hanseniaspora uvarum*, *Candida intermedia*, *C. crusei* (*Issatchenkia orientalis*), *C. parapsilosis*, *Brettanomyces bruxellensis*, *B. naardenensis*, *Dekkera anomala*, *Pichia membranaefaciens*, *P. minuta*, *P. anomala*, *Saccharomyces exiguus*, *S. cerevisiae*, *Saccharomycodes ludwigii* or for the specific detection of drink-spoiling molds of the genera *Mucor*, *Byssoschlamys*, *Neosartorya*, *Aspergillus* and *Talaromyces* in particular species of *Mucor racemosus*, *Byssoschlamys nivea*, *Neosartorya fischeri*, *Aspergillus fumigatus* and *A. fischeri*, *Talaromyces flavus*, *T. bacillisporus* and *T. flavus* or for the specific detection of drink-spoiling bacteria of the genera *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Weissella*, *Lactococcus*, *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter*, *Bacillus* and *Alicyclobacillus*, in particular species of *Lactobacillus collinoides*, *Leuconostoc mesenteroides*, *L. pseudomesenteroides*, *Oenococcus oeni*, *Bacillus coagulans*, *Alicyclobacillus ssp.*, *A. acidoterrestris*, *A. cycloheptanicus* and *A. herbarius* thus comprises the following steps:

- 25 - cultivating the drink-spoiling microorganisms present in the sample to be analysed
- fixing the drink-spoiling microorganisms present in the sample
- incubating the fixed drink-spoiling microorganisms with at least one nucleic acid probe and optionally in combination with a competitor probe, in order to achieve hybridization,
- 30 - removing or washing off the non-hybridized nucleic acid probe and

- detecting the drink-spoiling microorganisms hybridized to the nucleic acid probe molecules.

Within the present invention "cultivation" is understood to mean the propagation of  
5 the microorganisms present in the sample in a suitable cultivation medium.

For the detection of *yeasts and molds* the cultivation may occur, for example, in SSL-bouillon for 24 hours at 25°C. For the detection of lactic acid bacteria the cultivation may occur, for example, in MRS-bouillon for 48 hours at 30°C. For the  
10 detection of acetic acid bacteria the cultivation may occur, for example, on DSM-agar for 48 hours at 28°C. For the detection of bacilli, in particular *B. coagulans*, the cultivation may occur, for example, on dextrose-caseine-peptone agar for 48 hours at 55°C. For the detection of alicyclobacilli the cultivation may occur, for example, in BAM-bouillon for 48 hours at 44°C.

15

In any case, the person skilled in the art can find suitable cultivation methods in the prior art for each microorganism and each group of microorganisms to be analysed, respectively.

20 Within the present invention "fixing" of the microorganism is understood as a treatment with which the envelope of the microorganism is made permeable for nucleic acid probes. For fixation, usually ethanol is used. If the cell wall cannot be penetrated by the nucleic acid probes despite of using these techniques, the person skilled in the art will know enough further techniques which lead to the same result.  
25 These include, for example, methanol, mixtures of alcohols, low percentage paraformaldehyde solution or a diluted formaldehyde solution, enzymatic treatments or the like.

In a particularly preferred embodiment of the method of the present invention an  
30 enzymatic step may follow in order to cause complete cell disintegration of the

microorganisms. Enzymes which can accordingly be used for this step, are, for instance, lysozyme, proteinase K, and mutanolysine. The one skilled in the art will know sufficient suitable techniques and will easily find out which means is particularly suitable for cell disintegration of a certain microorganism.

5

Within the present invention the fixed microorganisms are incubated with fluorescently labeled nucleic acid probes for the "hybridization". These nucleic acid probes can, after the fixing, penetrate the cell wall and bind to the target sequence in the cell corresponding to the nucleic acid probe. Binding is to be understood as formation of hydrogen bonds between complementary nucleic acid pieces.

In such case the nucleic acid probe can be complementary to a chromosomal or episomal DNA, but can also be complementary to an mRNA or rRNA of the microorganism to be detected. It is advantageous to select a nucleic acid probe which is complementary to a region present in copies of more than 1 in the microorganism to be detected. The sequence to be detected is preferably present in 500-100,000 copies per cell, especially preferred 1,000-50,000 copies. For this reason the sequence of the rRNA is preferably used as a target site, since the ribosomes as sites of protein biosynthesis are present many thousandfold in each active cell.

The nucleic acid probe within the meaning of the invention may be a DNA or RNA probe comprising usually between 12 and 100 nucleotides, preferably between 15 and 50, more between 17 and 25 nucleotides. The selection of the nucleic acid probes is performed taking into consideration whether a complementary sequence is present in the microorganism to be detected. By this selection of a defined sequence, a species of a microorganism, a genus of a microorganism or an entire microorganism group may be detected. In a probe consisting of 15 nucleotides, the sequences should be 100% complementary. In case of oligonucleotides of more than 15 nucleotides, depending on the length of the oligonucleotide, one or more mismatches are allowed.

To increase the specificity of nucleic acid probes competitor probes can be used. Within the present invention competitor probes are understood to mean in particular oligonucleotides which block possibly undesired bindings of the nucleic acid probes and thereby show a higher sequence similarity to the non-target genera and species of microorganisms, respectively, than to the target genera and species of microorganisms, respectively. By using competitor probes the binding of the nucleic acid probe to the nucleic acid sequence of non-target genera or species of microorganisms can be prevented and thus does not lead to false signals. The non-labelled competitor probe is always used in combination with the labelled oligonucleotide probe.

The competitor probe should be complementary to a nucleic acid sequence having high sequence similarity to the nucleic acid sequence of the genera and species of microorganism, respectively, to be detected. In a particularly preferred embodiment the competitor probe is complementary to the rRNA of non-target genera and species of microorganism, respectively.

Within the meaning of the invention the competitor probe is a DNA or RNA sequence usually comprising between 12 and 100 nucleotides, preferably between 15 and 50, particularly preferably between 17 and 25 nucleotides. By selecting a defined sequence, a bacterial species, a bacterial genus or an entire bacterial group may be blocked. A probe consisting of 15 nucleotides should be 100% complementary to the nucleic acid sequence to be blocked. In case of oligonucleotides consisting of more than 15 nucleotides, depending on the length of the oligonucleotide, one or more mismatches are allowed.

Within the methods of the present invention the nucleic acid probe molecules of the present invention have the following lengths and sequences (all nucleic acid probe molecules are noted in 5'-3' direction).

The nucleic acid probe molecules of the present invention are useful for the specific detection of drink-spoiling yeasts of the genera *Zygosaccharomyces*, *Hanseniaspora*, *Candida*, *Brettanomyces*, *Dekkera*, *Pichia*, *Saccharomyces* and *Saccharomycodes* in particular the species *Zygosaccharomyces bailii*, *Z. mellis*, *Z. rouxii*, *Z. bisporus*, *Z. fermentati*, *Z. microellipsoides*, *Hanseniaspora uvarum*, *Candida intermedia*, *C. crusei* (*Issatchenkia orientalis*), *C. parapsilosis*, *Brettanomyces bruxellensis*, *B. naardenensis*, *Dekkera anomala*, *Pichia membranaefaciens*, *P. minuta*, *P. anomala*, *Saccharomyces exiguus*, *S. cerevisiae*, *Saccharomycodes ludwigii* or for the specific detection of drink-spoiling molds of the genera *Mucor*, *Byssoschlamys*, *Neosartorya*, *Aspergillus* and *Talaromyces* in particular species of *Mucor racemosus*, *Byssoschlamys nivea*, *Neosartorya fischeri*, *Aspergillus fumigatus* and *A. fischeri*, *Talaromyces flavus*, *T. bacillisporus* and *T. flavus* or for the specific detection of drink-spoiling bacteria of the genera *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Weissella*, *Lactococcus*, *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter*, *Bacillus* and *Alicyclobacillus*, in particular species of *Lactobacillus collinoides*, *Leuconostoc mesenteroides*, *L. pseudomesenteroides*, *Oenococcus oeni*, *Bacillus coagulans*, *Alicyclobacillus ssp.*, *A. acidoterrestris*, *A. cycloheptanicus* and *A. herbarius* and are used correspondingly in the detection method according to the invention.

Within the present invention probes that detect different species of microorganisms can be used in combination, in order to enable the simultaneous detection of different microorganisms. This leads likewise to an increase of speed of the detection method.

a) Nucleic acid molecules which specifically detect drink-spoiling yeasts:

SEQ ID No. 1: 5'-GTTTGACCAGATTCTCCGCTC

The sequence SEQ ID No. 1 is particularly useful for the detection of microorganisms of the genus *Zygosaccharomyces*.

SEQ ID No. 2: 5'- GTTTGACCAGATTTTCCGCTCT  
SEQ ID No. 3: 5'- GTTTGACCAAATTTTCCGCTCT  
SEQ ID No. 4: 5'- GTTTGTCCAAATTCTCCGCTCT

5

The nucleic acid molecules according to SEQ ID No. 2 to SEQ ID No. 4 are used as unlabelled competitor probes for the detection of microorganisms of the genus *Zygosaccharomyces* in combination with the nucleic acid probe according to SEQ ID No. 1 in order to prevent the binding of the labelled nucleic acid probe specific for members of the genus *Zygosaccharomyces* to nucleic acid sequences, which are not specific for members of the genus *Zygosaccharomyces*.

15 SEQ ID No. 5: 5'- CCCGGTCGAATTAAAACC  
SEQ ID No. 6: 5'- GCCCGGTCGAATTAAAAC  
SEQ ID No. 7: 5'- GGCCCGGTCGAATTAAAA  
SEQ ID No. 8: 5'- AGGCCCGGTCGAATTAAA  
SEQ ID No. 9: 5'- AAGGCCCGGTCGAATTAA  
SEQ ID No. 10: 5'- ATATTCGAGCGAAACGCC  
SEQ ID No. 11: 5'- AAAGATCCGGACCGGCCG  
20 SEQ ID No. 12 5'- GGAAAGATCCGGACCGGC  
SEQ ID No. 13 5'- GAAAGATCCGGACCGGCC  
SEQ ID No. 14 5'- GATCCGGACCGGCCGACC  
SEQ ID No. 15 5'- AGATCCGGACCGGCCGAC  
SEQ ID No. 16 5'- AAGATCCGGACCGGCCGA  
25 SEQ ID No. 17 5'- GAAAGGCCCGGTCGAATT  
SEQ ID No. 18 5'- AAAGGCCCGGTCGAATTA  
SEQ ID No. 19 5'- GGAAAGGCCCGGTCGAAT  
SEQ ID No. 20 5'- AGGAAAGGCCCGGTCGAA  
SEQ ID No. 21 5'- AAGGAAAGGCCCGGTCGA

30

The sequences SEQ ID No. 5 to SEQ ID No. 21 are particularly suitable for the detection of *Zygosaccharomyces bailii*.

SEQ ID No. 22: 5'- ATAGCACTGGGATCCTCGCC

5

The sequence SEQ ID No. 22 is particularly suitable for the detection of *Zygosaccharomyces fermentati*.

SEQ ID No. 23: 5'- CCAGCCCCAAAGTTACCTTC

10 SEQ ID No. 24: 5'- TCCTTGACGTAAAGTCGCAG

The sequences SEQ ID No. 23 to SEQ ID No. 24 are particularly suitable for the detection of *Zygosaccharomyces microellipsoides*.

15 SEQ ID No. 25: 5'- GGAAGAAAACCAGTACGC

SEQ ID No. 26: 5'- CCGGTCGGAAGAAAACCA

SEQ ID No. 27: 5'- GAAGAAAACCAGTACGCG

SEQ ID No. 28: 5'- CCCGGTCGGAAGAAAACC

SEQ ID No. 29: 5'- CGGTCGGAAGAAAACCAG

20 SEQ ID No. 30: 5'- GGTCGGAAGAAAACCAGT

SEQ ID No. 31: 5'- AAGAAAACCAGTACGCGG

SEQ ID No. 32: 5'- GTACGCGGAAAAATCCGG

SEQ ID No. 33: 5'- AGTACGCGGAAAAATCCG

SEQ ID No. 34: 5'- GCGGAAAAATCCGGACCG

25 SEQ ID No. 35: 5'- CGGAAGAAAACCAGTACG

SEQ ID No. 36: 5'- GCCCGGTCGGAAGAAAAC

SEQ ID No. 37: 5'- CGCGGAAAAATCCGGACC

SEQ ID No. 38: 5'- CAGTACGCGGAAAAATCC

SEQ ID No. 39: 5'- AGAAAACCAGTACGCGGA

30 SEQ ID No. 40: 5'- GGCCCGGTCGGAAGAAAA

	SEQ ID No. 41:	5'- ATAAACACCAACCCGATCC
	SEQ ID No. 42:	5'- ACGCGGAAAAATCCGGAC
	SEQ ID No. 43:	5'- GAGAGGCCCGGTCGGAAG
	SEQ ID No. 44:	5'- AGAGGCCCGGTCGGAAGA
5	SEQ ID No. 45:	5'- GAGGCCCGGTCGGAAGAA
	SEQ ID No. 46:	5'- AGGCCCGGTCGGAAGAAA
	SEQ ID No. 47:	5'- CCGAGTGGGTCAGTAAAT
	SEQ ID No. 48:	5'- CCAGTACGCGGAAAAATC
	SEQ ID No. 49:	5'- TAAACACCAACCCGATCCC
10	SEQ ID No. 50:	5'- GGAGAGGCCCGGTCGGA
	SEQ ID No. 51:	5'- GAAAACCAGTACGCGGAA
	SEQ ID No. 52:	5'- TACGCGGAAAAATCCGGA
	SEQ ID No. 53:	5'- GGCCACAGGGACCCAGGG
	SEQ ID No. 54:	5'- TCACCAAGGGCCACAGGG
15	SEQ ID No. 55:	5'- GGGCCACAGGGACCCAGG
	SEQ ID No. 56:	5'- TTCACCAAGGGCCACAGG
	SEQ ID No. 57:	5'- ACAGGGACCCAGGGCTAG
	SEQ ID No. 58:	5'- AGGGCCACAGGGACCCAG
	SEQ ID No. 59:	5'- GTTCACCAAGGGCCACAG
20	SEQ ID No. 60:	5'- GCCACAGGGACCCAGGGC
	SEQ ID No. 61:	5'- CAGGGACCCAGGGCTAGC
	SEQ ID No. 62:	5'- AGGGACCCAGGGCTAGCC
	SEQ ID No. 63:	5'- ACCAAGGGCCACAGGGAC
	SEQ ID No. 64:	5'- CCACAGGGACCCAGGGCT
25	SEQ ID No. 65:	5'- CACAGGGACCCAGGGCTA
	SEQ ID No. 66:	5'- CACCAAGGGCCACAGGGA
	SEQ ID No. 67:	5'- GGGACCCAGGGCTAGCCA
	SEQ ID No. 68:	5'- AGGAGAGGCCCGGTCGGA
	SEQ ID No. 69:	5'- AAGGAGAGGCCCGGTCGG
30	SEQ ID No. 70:	5'- GAAGGAGAGGCCCGGTCG



SEQ ID No. 71: 5'- AGGGCTAGCCAGAAGGAG  
SEQ ID No. 72: 5'- GGGCTAGCCAGAAGGAGA  
SEQ ID No. 73: 5'- AGAAGGAGAGGCCCGGTC  
SEQ ID No. 74: 5'- CAAGGGCCACAGGGACCC  
5 SEQ ID No. 75: 5'- CCAAGGGCCACAGGGACC

The sequences SEQ ID No. 25 to SEQ ID No. 75 are particularly suitable for the detection of *Zygosaccharomyces mellis*.

10 SEQ ID No. 76: 5'- GTCGGAAAAACCAGTACG  
SEQ ID No. 77: 5'- GCCCGGTCGGAAAAACCA  
SEQ ID No. 78: 5'- CCGGTCGGAAAAACCAGT  
SEQ ID No. 79: 5'- CCCGGTCGGAAAAACCAG  
SEQ ID No. 80: 5'- TCGGAAAAACCAGTACGC  
15 SEQ ID No. 81: 5'- CGGAAAAACCAGTACGCG  
SEQ ID No. 82: 5'- GGAAAAACCAGTACGCGG  
SEQ ID No. 83: 5'- GTACGCGGAAAAATCCGG  
SEQ ID No. 84: 5'- AGTACGCGGAAAAATCCG  
SEQ ID No. 85: 5'- GCGGAAAAATCCGGACCG  
20 SEQ ID No. 86: 5'- GGTCGGAAAAACCAGTAC  
SEQ ID No. 87: 5'- ACTCCTAGTGGTGCCCTT  
SEQ ID No. 88: 5'- GCTCCACTCCTAGTGGTG  
SEQ ID No. 89: 5'- CACTCCTAGTGGTGCCCT  
SEQ ID No. 90: 5'- CTCCACTCCTAGTGGTG  
25 SEQ ID No. 91: 5'- TCCACTCCTAGTGGTGCC  
SEQ ID No. 92: 5'- CCACTCCTAGTGGTGCCC  
SEQ ID No. 93: 5'- GGCTCCACTCCTAGTGGT  
SEQ ID No. 94: 5'- AGGCTCCACTCCTAGTGG  
SEQ ID No. 95: 5'- GGCCCGGTCGGAAAAACC  
30 SEQ ID No. 96: 5'- GAAAAACCAGTACGCGGA

	SEQ ID No. 97:	5'- CGCGGAAAAATCCGGACC
	SEQ ID No. 98:	5'- CAGTACGCGGAAAAATCC
	SEQ ID No. 99:	5'- CGGTCGGAAAAACAGTA
	SEQ ID No. 100:	5'- AAGGCCCGGTTCGGAAAAA
5	SEQ ID No. 101:	5'- CAGGCTCCACTCCTAGTG
	SEQ ID No. 102:	5'- CTCCTAGTGGTGCCCTTC
	SEQ ID No. 103:	5'- TCCTAGTGGTGCCCTTCC
	SEQ ID No. 104:	5'- GCAGGCTCCACTCCTAGT
	SEQ ID No. 105:	5'- AGGCCCGGTTCGGAAAAAC
10	SEQ ID No. 106:	5'- ACGCGGAAAAATCCGGAC
	SEQ ID No. 107:	5'- CCAGTACGCGGAAAAATC
	SEQ ID No. 108:	5'- CTAGTGGTGCCCTTCCGT
	SEQ ID No. 109:	5'- GAAAGGCCCGGTTCGGAAA
	SEQ ID No. 110:	5'- AAAGGCCCGGTTCGGAAAA
15	SEQ ID No. 111:	5'- TACGCGGAAAAATCCGGA
	SEQ ID No. 112:	5'- GGAAAGGCCCGGTTCGGAA
	SEQ ID No. 113:	5'- ATCTCTTCCGAAAGGTCG
	SEQ ID No. 114:	5'- CATCTCTTCCGAAAGGTC
	SEQ ID No. 115:	5'- CTCTTCCGAAAGGTCGAG
20	SEQ ID No. 116:	5'- CTTCCGAAAGGTCGAGAT
	SEQ ID No. 117:	5'- TCTCTTCCGAAAGGTCGA
	SEQ ID No. 118:	5'- TCTTCCGAAAGGTCGAGA
	SEQ ID No. 119:	5'- CCTAGTGGTGCCCTTCCG
	SEQ ID No. 120:	5'- TAGTGGTGCCCTTCCGTC
25	SEQ ID No. 121:	5'- AGTGGTGCCCTTCCGTCA
	SEQ ID No. 122:	5'- GCCAAGGTTAGACTCGTT
	SEQ ID No. 123:	5'- GGCCAAGGTTAGACTCGT
	SEQ ID No. 124:	5'- CCAAGGTTAGACTCGTTG
	SEQ ID No. 125:	5'- CAAGGTTAGACTCGTTGG
30	SEQ ID No. 126:	5'- AAGGTTAGACTCGTTGGC

The sequences SEQ ID No. 76 to SEQ ID No. 126 are particularly suitable for the detection of *Zygosaccharomyces rouxii*.

5 SEQ ID No. 127: 5'- CTCGCCTCACGGGGTTCTCA

The sequence SEQ ID No. 127 is particularly suitable for the simultaneous detection of *Zygosaccharomyces mellis* and *Zygosaccharomyces rouxii*.

10 SEQ ID No. 128: 5'- GGCCCGGTCGAAATTAAA  
SEQ ID No. 129: 5'- AGGCCCGGTCGAAATTAA  
SEQ ID No. 130: 5'- AAGGCCCGGTCGAAATTA  
SEQ ID No. 131: 5'- AAAGGCCCGGTCGAAATT  
SEQ ID No. 132: 5'- GAAAGGCCCGGTCGAAAT  
15 SEQ ID No. 133: 5'- ATATTCGAGCGAAACGCC  
SEQ ID No. 134: 5'- GGAAAGGCCCGGTCGAAA  
SEQ ID No. 135: 5'- AAAGATCCGGACCGGCCG  
SEQ ID No. 136: 5'- GGAAAGATCCGGACCGGC  
SEQ ID No. 137: 5'- GAAAGATCCGGACCGGCC  
20 SEQ ID No. 138: 5'- GATCCGGACCGGCCGACC  
SEQ ID No. 139: 5'- AGATCCGGACCGGCCGAC  
SEQ ID No. 140: 5'- AAGATCCGGACCGGCCGA  
SEQ ID No. 141: 5'- AGGAAAGGCCCGGTCGAA  
SEQ ID No. 142: 5'- AAGGAAAGGCCCGGTCGA

25

The sequences SEQ ID No. 128 to SEQ ID No. 142 are particularly suitable for the detection of *Zygosaccharomyces bisporus*.

SEQ ID No. 143: 5'-CGAGCAAAACGCCTGCTTTG  
30 SEQ ID No. 144: 5'-CGCTCTGAAAGAGAGTTGCC

The sequences SEQ ID No. 143 and SEQ ID No. 144 are particularly suitable for the detection of *Hanseniaspora uvarum*.

5    SEQ ID No. 145:    5'-AGTTGCCCCCTACACTAGAC  
    SEQ ID No. 146:    5'-GCTTCTCCGTCCCGCGCCG

The sequences SEQ ID No. 145 and SEQ ID No. 146 are particularly suitable for the detection of *Candida intermedia*.

10

SEQ ID No. 147:    5'- AGATTYTCCGCTCTGAGATGG

The nucleic acid probe molecule according to SEQ ID No. 147 is used as unlabelled competitor probe for the detection of *Candida intermedia* in combination with the  
15    oligonucleotide probe according to SEQ ID No. 146, in order to prevent the binding of the labelled oligonucleotide probe specific for *Candida intermedia* to nucleic acid sequences which are not specific for *Candida intermedia*.

20    SEQ ID No. 148:    5'- CCTGGTTCGCCAAAAAGGC

The sequence SEQ ID No. 148 is particularly suitable for the detection of *Candida parapsilosis*.

25    SEQ ID No. 149:    5'-GATTCTCGGCCCCATGGG

The sequence SEQ ID No. 149 is particularly suitable for the detection of *Candida crusei* (*Issatchenkia orientalis*).

30    SEQ ID No. 150:    5'- ACCCTCTACGGCAGCCTGTT

The sequence SEQ ID No. 150 is particularly suitable for the detection of *Dekkera anomala* and *Brettanomyces (Dekkera) bruxellensis*.

SEQ ID No. 151: 5'- GATCGGTCTCCAGCGATTCA

5

The sequence SEQ ID No. 151 is particularly suitable for the detection of *Brettanomyces (Dekkera) bruxellensis*.

SEQ ID No. 152: 5'- ACCCTCCACGGCGGCCTGTT

10

The sequence SEQ ID No. 152 is particularly suitable for the detection of *Brettanomyces (Dekkera) naardenensis*.

SEQ ID No. 153: 5'- GATTCTCCGCGCCATGGG

15

The sequence SEQ ID No. 153 is particularly suitable for the detection of *Pichia membranaefaciens*.

SEQ ID No. 154: 5'- TCATCAGACGGGATTCTCAC

20

The sequence SEQ ID No. 154 is particularly suitable for the simultaneous detection of *Pichia minuta* and *Pichia anomala*.

SEQ ID No. 155: 5'- CTCATCGCACGGGATTCTCACC

25

SEQ ID No. 156: 5'- CTCGCCACACGGGATTCTCACC

The nucleic acid probe molecules according to SEQ ID No. 155 and SEQ ID No. 156 are used as unlabelled competitor probes for the simultaneous detection of *Pichia minuta* and *Pichia anomala* in combination with the oligonucleotide probe according to SEQ ID No. 154, in order to prevent the binding of the labelled oligonucleotide

30

probe specific for *Pichia minuta* and *Pichia anomala*, to nucleic acid sequences which are not specific for *Pichia minuta* and *Pichia anomala*.

SEQ ID No. 157: 5'-AGTTGCCCCCTCCTCTAAGC

5

The sequence SEQ ID No. 157 is particularly suitable for the detection of *Saccharomyces exiguus*.

SEQ ID No. 158: 5'-CTGCCACAAGGACAAATGGT

10 SEQ ID No. 159: 5'-TGCCCCCTCTTCTAAGCAAAT

The sequences SEQ ID No. 158 and SEQ ID No. 159 are particularly suitable for the detection of *Saccharomyces ludwigii*.

15 SEQ ID No. 160: 5'-CCCCAAAGTTGCCCTCTC

The sequence SEQ ID No. 160 is particularly suitable for the detection of *Saccharomyces cerevisiae*.

20 SEQ ID No. 161: 5'-GCCGCCCCAAAGTCGCCCTCTAC

SEQ ID No. 162: 5'-GCCCCAGAGTCGCCTTCTAC

The nucleic acid probe molecules according to SEQ ID No. 161 and SEQ ID No. 162 are used as unlabelled competitor probes for the detection of *Saccharomyces cerevisiae* in combination with the oligonucleotide probe according to SEQ ID No. 160, in order to prevent the binding of the labelled oligonucleotide probe specific for *Saccharomyces cerevisiae*, to nucleic acid sequences which are not specific for *Saccharomyces cerevisiae*.

30 b) Nucleic acid probe molecules which specifically detect drink-spoiling molds:

SEQ ID No. 163: 5'-AAGACCAGGCCACCTCAT

5 The sequence SEQ ID No. 163 is particularly suitable for the detection of *Mucor racemosus*.

SEQ ID No. 164: 5'-CATCATAGAACACCGTCC

10 The sequence SEQ ID No. 164 is particularly suitable for the detection of *Byssoschlamys nivea*.

SEQ ID No. 165: 5'-CCTTCCGAAGTCGAGGTTTT

15 The sequence SEQ ID No. 165 is particularly suitable for the detection of *Neosartorya fischeri*.

SEQ ID No. 166: 5'-GGGAGTGTTGCCAACTC

20 The sequence SEQ ID No. 166 is particularly suitable for the simultaneous detection of *Aspergillus fumigatus* and *A. fischeri*.

SEQ ID No. 167: 5'-AGCGGTCGTTCGCAACCCT

25 The sequence SEQ ID No. 167 is particularly suitable for the detection of *Talaromyces flavus*.

SEQ ID No. 168: 5'-CCGAAGTCGGGGTTTTGCGG

30 The sequence SEQ ID No. 168 is particularly suitable for the simultaneous detection of *Talaromyces bacillisporus* and *T. flavus*.

c) Nucleic acid probe molecules, which specifically detect drink-spoiling lactic acid bacteria

5	SEQ ID No. 169:	5'- GATAGCCGAAACCACCTTTC
	SEQ ID No. 170:	5'- GCCGAAACCACCTTTCAAAC
	SEQ ID No. 171:	5'- GTGATAGCCGAAACCACCTT
	SEQ ID No. 172:	5'- AGTGATAGCCGAAACCACCT
	SEQ ID No. 173:	5'- TTTAACGGGATGCGTTCGAC
10	SEQ ID No. 174:	5'- AAGTGATAGCCGAAACCACC
	SEQ ID No. 175:	5'- GGTTGAATACCGTCAACGTC
	SEQ ID No. 176:	5'- GCACAGTATGTCAAGACCTG
	SEQ ID No. 177:	5'- CATCCGATGTGCAAGCACTT
	SEQ ID No. 178:	5'- TCATCCGATGTGCAAGCACT
15	SEQ ID No. 179:	5'- CCGATGTGCAAGCACTTCAT
	SEQ ID No. 180:	5'- CCACTCATCCGATGTGCAAG
	SEQ ID No. 181:	5'- GCCACAGTTCGCCACTCATC
	SEQ ID No. 182:	5'- CCTCCGCGTTTGTACACGGC
	SEQ ID No. 183:	5'- ACCAGTTCGCCACAGTTCGC
20	SEQ ID No. 184:	5'- CACTCATCCGATGTGCAAGC
	SEQ ID No. 185:	5'- CCAGTTCGCCACAGTTCGCC
	SEQ ID No. 186:	5'- CTCATCCGATGTGCAAGCAC
	SEQ ID No. 187:	5'- TCCGATGTGCAAGCACTTCA
	SEQ ID No. 188:	5'- CGCCACTCATCCGATGTGCA
25	SEQ ID No. 189:	5'- CAGTTCGCCACAGTTCGCCA
	SEQ ID No. 190:	5'- GCCACTCATCCGATGTGCAA
	SEQ ID No. 191:	5'- CGCCACAGTTCGCCACTCAT
	SEQ ID No. 192:	5'- ATCCGATGTGCAAGCACTTC
	SEQ ID No. 193:	5'- GTTCGCCACAGTTCGCCACT
30	SEQ ID No. 194:	5'- TCCTCCGCGTTTGTACACGG



SEQ ID No. 195: 5'- CGCCAGGGTTCATCCTGAGC  
SEQ ID No. 196: 5'- AGTTCGCCACAGTTCGCCAC  
SEQ ID No. 197: 5'- TCGCCACAGTTCGCCACTCA  
SEQ ID No. 198: 5'- TTAACGGGATGCGTTCGACT  
5 SEQ ID No. 199: 5'- TCGCCACTCATCCGATGTGC  
SEQ ID No. 200: 5'- CCACAGTTCGCCACTCATCC  
SEQ ID No. 201: 5'- GATTTAACGGGATGCGTTTCG  
SEQ ID No. 202: 5'- TAACGGGATGCGTTCGACTT  
SEQ ID No. 203: 5'- AACGGGATGCGTTCGACTTG  
10 SEQ ID No. 204: 5'- CGAAGGTTACCGAACCGACT  
SEQ ID No. 205: 5'- CCGAAGGTTACCGAACCGAC  
SEQ ID No. 206: 5'- CCCGAAGGTTACCGAACCGA  
SEQ ID No. 207: 5'- TTCCTCCGCGTTTGTCAACG  
SEQ ID No. 208: 5'- CCGCCAGGGTTCATCCTGAG  
15 SEQ ID No. 209: 5'- TCCTTCCAGAAGTGATAGCC  
SEQ ID No. 210: 5'- CACCAGTTCGCCACAGTTTCG  
SEQ ID No. 211: 5'- ACGGGATGCGTTCGACTTGC  
SEQ ID No. 212: 5'- GTCCTTCCAGAAGTGATAGC  
SEQ ID No. 213: 5'- GCCAGGGTTCATCCTGAGCC  
20 SEQ ID No. 214: 5'- ACTCATCCGATGTGCAAGCA  
SEQ ID No. 215: 5'- ATCATTGCCTTGGTGAACCG  
SEQ ID No. 216: 5'- TCCGCGTTTGTCAACCGGCAG  
SEQ ID No. 217: 5'- TGAACCGTTACTCCACCAAC  
SEQ ID No. 218: 5'- GAAGTGATAGCCGAAACCAC  
25 SEQ ID No. 219: 5'- CCGCGTTTGTCAACCGGCAGT  
SEQ ID No. 220: 5'- TTCGCCACTCATCCGATGTG  
SEQ ID No. 221: 5'- CATTTAACGGGATGCGTTTCG  
SEQ ID No. 222: 5'- CACAGTTCGCCACTCATCCG  
SEQ ID No. 223: 5'- TTCGCCACAGTTCGCCACTC  
30 SEQ ID No. 224: 5'- CTCCGCGTTTGTCAACCGGCA

	SEQ ID No. 225:	5'- ACGCCGCCAGGGTTCATCCT
	SEQ ID No. 226:	5'- CCTTCCAGAAGTGATAGCCG
	SEQ ID No. 227:	5'- TCATTGCCTTGGTGAACCGT
	SEQ ID No. 228:	5'- CACAGTATGTCAAGACCTGG
5	SEQ ID No. 229:	5'- TTGGTGAACCGTTACTCCAC
	SEQ ID No. 230:	5'- CTTGGTGAACCGTTACTCCA
	SEQ ID No. 231:	5'- GTGAACCGTTACTCCACCAA
	SEQ ID No. 232:	5'- GGCTCCCGAAGGTTACCGAA
	SEQ ID No. 233:	5'- GAAGGTTACCGAACCGACTT
10	SEQ ID No. 234:	5'- TGGCTCCCGAAGGTTACCGA
	SEQ ID No. 235:	5'- TAATACGCCGCGGGTCCTTC
	SEQ ID No. 236:	5'- GAACCGTTACTCCACCAACT
	SEQ ID No. 237:	5'- TACGCCGCGGGTCCTTCCAG
	SEQ ID No. 238:	5'- TCACCAGTTCGCCACAGTTC
15	SEQ ID No. 239:	5'- CCTTGGTGAACCGTTACTCC
	SEQ ID No. 240:	5'- CTCACCAGTTCGCCACAGTT
	SEQ ID No. 241:	5'- CGCCGCCAGGGTTCATCCTG
	SEQ ID No. 242:	5'- CCTTGGTGAACCATTACTCC
	SEQ ID No. 243:	5'- TGGTGAACCATTACTCCACC
20	SEQ ID No. 244:	5'- GCCGCCAGGGTTCATCCTGA
	SEQ ID No. 245:	5'- GGTGAACCATTACTCCACCA
	SEQ ID No. 246:	5'- CCAGGGTTCATCCTGAGCCA
	SEQ ID No. 247:	5'- AATACGCCGCGGGTCCTTCC
	SEQ ID No. 248:	5'- CACGCCGCCAGGGTTCATCC
25	SEQ ID No. 249:	5'- AGTTCGCCACTCATCCGATG
	SEQ ID No. 250:	5'- CGGGATGCGTTCGACTTGCA
	SEQ ID No. 251:	5'- CATTGCCTTGGTGAACCGTT
	SEQ ID No. 252:	5'- GCACGCCGCCAGGGTTCATC
	SEQ ID No. 253:	5'- CTCCTCCGCGTTTGTCACC
30	SEQ ID No. 254:	5'- TGGTGAACCGTTACTCCACC

SEQ ID No. 255: 5'- CCTTCCTCCGCGTTTGTAC  
SEQ ID No. 256: 5'- ACGCCGCGGGTCCTTCCAGA  
SEQ ID No. 257: 5'- GGTGAACCGTTACTCCACCA  
SEQ ID No. 258: 5'- GGGTCCTTCCAGAAGTGATA  
5 SEQ ID No. 259: 5'- CTTCCAGAAGTGATAGCCGA  
SEQ ID No. 260: 5'- GCCTTGGTGAACCATTACTC  
SEQ ID No. 261: 5'- ACAGTTCGCCACTCATCCGA  
SEQ ID No. 262: 5'- ACCTTCCTCCGCGTTTGTCA  
SEQ ID No. 263: 5'- CGAACCGACTTTGGGTGTTG  
10 SEQ ID No. 264: 5'- GAACCGACTTTGGGTGTTGC  
SEQ ID No. 265: 5'- AGGTTACCGAACCGACTTTG  
SEQ ID No. 266: 5'- ACCGAACCGACTTTGGGTGT  
SEQ ID No. 267: 5'- TTACCGAACCGACTTTGGGT  
SEQ ID No. 268: 5'- TACCGAACCGACTTTGGGTG  
15 SEQ ID No. 269: 5'- GTTACCGAACCGACTTTGGG

The sequences SEQ ID No. 169 to SEQ ID No. 269 are particularly suitable for the detection of *Lactobacillus collinoides*.

20 SEQ ID No. 270: 5'- CCTTTCTGGTATGGTACCGTC  
SEQ ID No. 271: 5'- TGCACCGCGGAYCCATCTCT

The sequences SEQ ID No. 270 to SEQ ID No. 271 are particularly suitable for the detection of members of the genus *Leuconostoc*.

25  
SEQ ID No. 272: 5'- AGTTGCAGTCCAGTAAGCCG  
SEQ ID No. 273: 5'- GTTGCAGTCCAGTAAGCCGC  
SEQ ID No. 274: 5'- CAGTTGCAGTCCAGTAAGCC  
SEQ ID No. 275: 5'- TGCAGTCCAGTAAGCCGCCT  
30 SEQ ID No. 276: 5'- TCAGTTGCAGTCCAGTAAGC

SEQ ID No. 277: 5'- TTGCAGTCCAGTAAGCCGCC  
SEQ ID No. 278: 5'- GCAGTCCAGTAAGCCGCCTT  
SEQ ID No. 279: 5'- GTCAGTTGCAGTCCAGTAAG  
SEQ ID No. 280: 5'- CTCTAGGTGACGCCGAAGCG  
5 SEQ ID No. 281: 5'- ATCTCTAGGTGACGCCGAAG  
SEQ ID No. 282: 5'- TCTAGGTGACGCCGAAGCGC  
SEQ ID No. 283: 5'- TCTCTAGGTGACGCCGAAGC  
SEQ ID No. 284: 5'- CCATCTCTAGGTGACGCCGA  
SEQ ID No. 285: 5'- CATCTCTAGGTGACGCCGAA  
10 SEQ ID No. 286: 5'- TAGGTGACGCCGAAGCGCCT  
SEQ ID No. 287: 5'- CTAGGTGACGCCGAAGCGCC  
SEQ ID No. 288: 5'- CTTAGACGGCTCCTTCCTAA  
SEQ ID No. 289: 5'- CCTTAGACGGCTCCTTCCTA  
SEQ ID No. 290: 5'- ACGTCAGTTGCAGTCCAGTA  
15 SEQ ID No. 291: 5'- CGTCAGTTGCAGTCCAGTAA  
SEQ ID No. 292: 5'- ACGCCGAAGCGCCTTTTAAC  
SEQ ID No. 293: 5'- GACGCCGAAGCGCCTTTTAA  
SEQ ID No. 294: 5'- GCCGAAGCGCCTTTTAACTT  
SEQ ID No. 295: 5'- CGCCGAAGCGCCTTTTAACT  
20 SEQ ID No. 296: 5'- GTGACGCCGAAGCGCCTTTT  
SEQ ID No. 297: 5'- TGACGCCGAAGCGCCTTTTA  
SEQ ID No. 298: 5'- AGACGGCTCCTTCCTAAAAG  
SEQ ID No. 299: 5'- ACGGCTCCTTCCTAAAAGGT  
SEQ ID No. 300: 5'- GACGGCTCCTTCCTAAAAGG  
25 SEQ ID No. 301: 5'- CCTTCCTAAAAGGTTAGGCC

The sequences SEQ ID No. 272 to SEQ ID No. 301 are particularly suitable for the simultaneous detection of *Leuconostoc mesenteroides* and *Leuconostoc pseudomesenteroides*.

SEQ ID No. 302: 5'- GGTGACGCCAAAGCGCCTTT  
SEQ ID No. 303: 5'- AGGTGACGCCAAAGCGCCTT  
SEQ ID No. 304: 5'- TAGGTGACGCCAAAGCGCCT  
SEQ ID No. 305: 5'- CTCTAGGTGACGCCAAAGCG  
5 SEQ ID No. 306: 5'- TCTAGGTGACGCCAAAGCGC  
SEQ ID No. 307: 5'- CTAGGTGACGCCAAAGCGCC  
SEQ ID No. 308: 5'- ACGCCAAAGCGCCTTTTAAC  
SEQ ID No. 309: 5'- CGCCAAAGCGCCTTTTAACT  
SEQ ID No. 310: 5'- TGACGCCAAAGCGCCTTTTA  
10 SEQ ID No. 311: 5'- TCTCTAGGTGACGCCAAAGC  
SEQ ID No. 312: 5'- GTGACGCCAAAGCGCCTTTT  
SEQ ID No. 313: 5'- GACGCCAAAGCGCCTTTTAA  
SEQ ID No. 314: 5'- ATCTCTAGGTGACGCCAAAG  
SEQ ID No. 315: 5'- CATCTCTAGGTGACGCCAAA  
15 SEQ ID No. 316: 5'- TCCATCTCTAGGTGACGCCA  
SEQ ID No. 317: 5'- CCATCTCTAGGTGACGCCAA  
SEQ ID No. 318: 5'- CTGCCTTAGACGGCTCCCCC  
SEQ ID No. 319: 5'- CCTGCCTTAGACGGCTCCCC  
SEQ ID No. 320: 5'- GTGTCATGCGACACTGAGTT  
20 SEQ ID No. 321: 5'- TGTGTCATGCGACACTGAGT  
SEQ ID No. 322: 5'- CTTTGTGTCATGCGACACTG  
SEQ ID No. 323: 5'- TTGTGTCATGCGACACTGAG  
SEQ ID No. 324: 5'- TGCCTTAGACGGCTCCCCCT  
SEQ ID No. 325: 5'- AGACGGCTCCCCCTAAAAGG  
25 SEQ ID No. 326: 5'- TAGACGGCTCCCCCTAAAAG  
SEQ ID No. 327: 5'- GCCTTAGACGGCTCCCCCTA  
SEQ ID No. 328: 5'- GCTCCCCCTAAAAGGTTAGG  
SEQ ID No. 329: 5'- GGCTCCCCCTAAAAGGTTAG  
SEQ ID No. 330: 5'- CTCCCCCTAAAAGGTTAGGC  
30 SEQ ID No. 331: 5'- TCCCCCTAAAAGGTTAGGCC

SEQ ID No. 332: 5'- CCCTAAAAGGTTAGGCCACC  
 SEQ ID No. 333: 5'- CCCCTAAAAGGTTAGGCCAC  
 SEQ ID No. 334: 5'- CGGCTCCCCCTAAAAGGTTA  
 SEQ ID No. 335: 5'- CCCCCTAAAAGGTTAGGCCA  
 5 SEQ ID No. 336: 5'- CTTAGACGGCTCCCCCTAAA  
 SEQ ID No. 337: 5'- TTAGACGGCTCCCCCTAAAA  
 SEQ ID No. 338: 5'- GGGTTCGCAACTCGTTGTAT  
 SEQ ID No. 339: 5'- CCTTAGACGGCTCCCCCTAA  
 SEQ ID No. 340: 5'- ACGGCTCCCCCTAAAAGGTT  
 10 SEQ ID No. 341: 5'- GACGGCTCCCCCTAAAAGGT

The sequences SEQ ID No. 302 to SEQ ID No. 341 are particularly suitable for the detection of *Leuconostoc pseudomesenteroides*.

15 SEQ ID No. 342: 5'- ACGCCGCAAGACCATCCTCT  
 SEQ ID No. 343: 5'- CTAATACGCCGCAAGACCAT  
 SEQ ID No. 344: 5'- TACGCCGCAAGACCATCCTC  
 SEQ ID No. 345: 5'- GTTACGATCTAGCAAGCCGC  
 SEQ ID No. 346: 5'- AATACGCCGCAAGACCATCC  
 20 SEQ ID No. 347: 5'- CGCCGCAAGACCATCCTCTA  
 SEQ ID No. 348: 5'- GCTAATACGCCGCAAGACCA  
 SEQ ID No. 349: 5'- ACCATCCTCTAGCGATCCAA  
 SEQ ID No. 350: 5'- TAATACGCCGCAAGACCATC  
 SEQ ID No. 351: 5'- AGCCATCCCTTTCTGGTAAG  
 25 SEQ ID No. 352: 5'- ATACGCCGCAAGACCATCCT  
 SEQ ID No. 353: 5'- AGTTACGATCTAGCAAGCCG  
 SEQ ID No. 354: 5'- AGCTAATACGCCGCAAGACC  
 SEQ ID No. 355: 5'- GCCGCAAGACCATCCTCTAG  
 SEQ ID No. 356: 5'- TTACGATCTAGCAAGCCGCT  
 30 SEQ ID No. 357: 5'- GACCATCCTCTAGCGATCCA

SEQ ID No. 358: 5'- TTGCTACGTCAGTAGGAGGC  
SEQ ID No. 359: 5'- ACGTCACTAGGAGGCGGAAA  
SEQ ID No. 360: 5'- TTTGCTACGTCAGTAGGAGG  
SEQ ID No. 361: 5'- GCCATCCCTTTCTGGTAAGG  
5 SEQ ID No. 362: 5'- TACGTCAGTAGGAGGCGGAA  
SEQ ID No. 363: 5'- CGTCACTAGGAGGCGGAAAC  
SEQ ID No. 364: 5'- AAGACCATCCTCTAGCGATC  
SEQ ID No. 365: 5'- GCACGTATTTAGCCATCCCT  
SEQ ID No. 366: 5'- CTCTAGCGATCCAAAAGGAC  
10 SEQ ID No. 367: 5'- CCTCTAGCGATCCAAAAGGA  
SEQ ID No. 368: 5'- CCATCCTCTAGCGATCCAAA  
SEQ ID No. 369: 5'- GGCACGTATTTAGCCATCCC  
SEQ ID No. 370: 5'- TACGATCTAGCAAGCCGCTT  
SEQ ID No. 371: 5'- CAGTTACGATCTAGCAAGCC  
15 SEQ ID No. 372: 5'- CCGCAAGACCATCCTCTAGC  
SEQ ID No. 373: 5'- CCATCCCTTTCTGGTAAGGT  
SEQ ID No. 374: 5'- AGACCATCCTCTAGCGATCC  
SEQ ID No. 375: 5'- CAAGACCATCCTCTAGCGAT  
SEQ ID No. 376: 5'- GCTACGTCAGTAGGAGGCGG  
20 SEQ ID No. 377: 5'- TGCTACGTCAGTAGGAGGCG  
SEQ ID No. 378: 5'- CTACGTCAGTAGGAGGCGGA  
SEQ ID No. 379: 5'- CCTCAACGTCAGTTACGATC  
SEQ ID No. 380: 5'- GTCAGTAGGAGGCGGAAACC  
SEQ ID No. 381: 5'- TCCTCTAGCGATCCAAAAGG  
25 SEQ ID No. 382: 5'- TGGCACGTATTTAGCCATCC  
SEQ ID No. 383: 5'- ACGATCTAGCAAGCCGCTTT  
SEQ ID No. 384: 5'- GCCAGTCTCTCAACTCGGCT  
SEQ ID No. 385: 5'- AAGCTAATACGCCGCAAGAC  
SEQ ID No. 386: 5'- GTTTGCTACGTCAGTAGGAG  
30 SEQ ID No. 387: 5'- CGCCACTCTAGTCATTGCCT

	SEQ ID No. 388:	5'- GGCCAGCCAGTCTCTCAACT
	SEQ ID No. 389:	5'- CAGCCAGTCTCTCAACTCGG
	SEQ ID No. 390:	5'- CCCGAAGATCAATTCAGCGG
	SEQ ID No. 391:	5'- CCGGCCAGTCTCTCAACTCG
5	SEQ ID No. 392:	5'- CCAGCCAGTCTCTCAACTCG
	SEQ ID No. 393:	5'- TCATTGCCTCACTTCACCCG
	SEQ ID No. 394:	5'- GCCAGCCAGTCTCTCAACTC
	SEQ ID No. 395:	5'- CACCCGAAGATCAATTCAGC
	SEQ ID No. 396:	5'- GTCATTGCCTCACTTCACCC
10	SEQ ID No. 397:	5'- CATTGCCTCACTTCACCCGA
	SEQ ID No. 398:	5'- ATTGCCTCACTTCACCCGAA
	SEQ ID No. 399:	5'- CGAAGATCAATTCAGCGGCT
	SEQ ID No. 400:	5'- AGTCATTGCCTCACTTCACC
	SEQ ID No. 401:	5'- TCGCCACTCTAGTCATTGCC
15	SEQ ID No. 402:	5'- TTGCCTCACTTCACCCGAAG
	SEQ ID No. 403:	5'- CGGCCAGTCTCTCAACTCGG
	SEQ ID No. 404:	5'- CTGGCACGTATTTAGCCATC
	SEQ ID No. 405:	5'- ACCCGAAGATCAATTCAGCG
	SEQ ID No. 406:	5'- TCTAGCGATCCAAAAGGACC
20	SEQ ID No. 407:	5'- CTAGCGATCCAAAAGGACCT
	SEQ ID No. 408:	5'- GCACCCATCGTTTACGGTAT
	SEQ ID No. 409:	5'- CACCCATCGTTTACGGTATG
	SEQ ID No. 410:	5'- GCCACTCTAGTCATTGCCTC
	SEQ ID No. 411:	5'- CGTTTGCTACGTCACTAGGA
25	SEQ ID No. 412:	5'- GCCTCAACGTCAGTTACGAT
	SEQ ID No. 413:	5'- GCCGGCCAGTCTCTCAACTC
	SEQ ID No. 414:	5'- TCACTAGGAGGCGGAAACCT
	SEQ ID No. 415:	5'- AGCCTCAACGTCAGTTACGA
	SEQ ID No. 416:	5'- AGCCAGTCTCTCAACTCGGC..
30	SEQ ID No. 417:	5'- GGCCAGTCTCTCAACTCGGC...



SEQ ID No. 418: 5'- CAAGCTAATACGCCGCAAGA  
SEQ ID No. 419: 5'- TTCGCCACTCTAGTCATTGC  
SEQ ID No. 420: 5'- CCGAAGATCAATTCAGCGGC  
SEQ ID No. 421: 5'- CGCAAGACCATCCTCTAGCG  
5 SEQ ID No. 422: 5'- GCAAGACCATCCTCTAGCGA  
SEQ ID No. 423: 5'- GCGTTTGCTACGTCACTAGG  
SEQ ID No. 424: 5'- CCACTCTAGTCATTGCCTCA  
SEQ ID No. 425: 5'- CACTCTAGTCATTGCCTCAC  
SEQ ID No. 426: 5'- CCAGTCTCTCAACTCGGCTA  
10 SEQ ID No. 427: 5'- TTACCTTAGGCACCGGCCTC  
SEQ ID No. 428: 5'- ACAAGCTAATACGCCGCAAG  
SEQ ID No. 429: 5'- TTTACCTTAGGCACCGGCCT  
SEQ ID No. 430: 5'- TTTTACCTTAGGCACCGGCC  
SEQ ID No. 431: 5'- ATTTTACCTTAGGCACCGGC  
15 SEQ ID No. 432: 5'- GATTTTACCTTAGGCACCGG  
SEQ ID No. 433: 5'- CTCACTTCACCCGAAGATCA  
SEQ ID No. 434: 5'- ACGCCACCAGCGTTCATCCT  
SEQ ID No. 435: 5'- GCCAAGCGACTTTGGGTACT  
SEQ ID No. 436: 5'- CGGAAAATTCCCTACTGCAG  
20 SEQ ID No. 437: 5'- CGATCTAGCAAGCCGCTTTC  
SEQ ID No. 438: 5'- GGTACCGTCAAGCTGAAAAC  
SEQ ID No. 439: 5'- TGCCTCACTTCACCCGAAGA  
SEQ ID No. 440: 5'- GGCCGGCCAGTCTCTCAACT  
SEQ ID No. 441: 5'- GGTAAGGTACCGTCAAGCTG  
25 SEQ ID No. 442: 5'- GTAAGGTACCGTCAAGCTGA  
SEQ ID No. 443: 5'- CCGCAAGACCATCCTCTAGG  
SEQ ID No. 444: 5'- ATTTAGCCATCCCTTTCTGG

The sequences SEQ ID No. 342 to SEQ ID No. 444 are particularly suitable for the  
30 detection of *Oenococcus oeni*.

SEQ ID No. 445: 5'- AACCTTCATCACACACG  
SEQ ID No. 446: 5'- CGAAACCCTTCATCACAC  
SEQ ID No. 447: 5'- ACCCTTCATCACACACGC  
5 SEQ ID No. 448: 5'- TACCGTCACACACTGAAC  
SEQ ID No. 449: 5'- AGATACCGTCACACACTG  
SEQ ID No. 450: 5'- CACTCAAGGGCGGAAACC  
SEQ ID No. 451: 5'- ACCGTCACACACTGAACA  
SEQ ID No. 452: 5'- CGTCACACACTGAACAGT  
10 SEQ ID No. 453: 5'- CCGAAACCCTTCATCACA  
SEQ ID No. 454: 5'- CCGTCACACACTGAACAG  
SEQ ID No. 455: 5'- GATACCGTCACACACTGA  
SEQ ID No. 456: 5'- GGTAAGATACCGTCACAC  
SEQ ID No. 457: 5'- CCCTTCATCACACACGCG  
15 SEQ ID No. 458: 5'- ACAGTGTTTTACGAGCCG  
SEQ ID No. 459: 5'- CAGTGTTTTACGAGCCGA  
SEQ ID No. 460: 5'- ACAAAGCGTTCGACTTGC  
SEQ ID No. 461: 5'- CGGATAACGCTTGGAACA  
SEQ ID No. 462: 5'- AGGGCGGAAACCCTCGAA  
20 SEQ ID No. 463: 5'- GGGCGGAAACCCTCGAAC  
SEQ ID No. 464: 5'- GGC GGAAACCCTCGAACA  
SEQ ID No. 465: 5'- TGAGGGCTTTCACCTCAG  
SEQ ID No. 466: 5'- AGGGCTTTCACCTCAGAC  
SEQ ID No. 467: 5'- GAGGGCTTTCACCTCAGA  
25 SEQ ID No. 468: 5'- ACTGCACTCAAGTCATCC  
SEQ ID No. 469: 5'- CCGGATAACGCTTGGAAC  
SEQ ID No. 470: 5'- TCCGGATAACGCTTGGA  
SEQ ID No. 471: 5'- TATCCCCTGCTAAGAGGT  
SEQ ID No. 472: 5'- CCTGCTAAGAGGTAGGTT  
30 SEQ ID No. 473: 5'- CCCTGCTAAGAGGTAGGT

SEQ ID No. 474: 5'- CCCCTGCTAAGAGGTAGG  
SEQ ID No. 475: 5'- TCCCCTGCTAAGAGGTAG  
SEQ ID No. 476: 5'- ATCCCCTGCTAAGAGGTA  
SEQ ID No. 477: 5'- CCGTTCCTTTCTGGTAAG  
5 SEQ ID No. 478: 5'- GCCGTTCCTTTCTGGTAA  
SEQ ID No. 479: 5'- AGCCGTTCCTTTCTGGTA  
SEQ ID No. 480: 5'- GCACGTATTTAGCCGTTT  
SEQ ID No. 481: 5'- CACGTATTTAGCCGTTCC  
SEQ ID No. 482: 5'- GGCACGTATTTAGCCGTT  
10 SEQ ID No. 483: 5'- CACTTTCCTCTACTGCAC  
SEQ ID No. 484: 5'- CCACTTTCCTCTACTGCA  
SEQ ID No. 485: 5'- TCCACTTTCCTCTACTGC  
SEQ ID No. 486: 5'- CTTTCCTCTACTGCACTC  
SEQ ID No. 487: 5'- TAGCCGTTTCCTTTCTGGT  
15 SEQ ID No. 488: 5'- TTAGCCGTTTCCTTTCTGG  
SEQ ID No. 489: 5'- TTATCCCCTGCTAAGAGG  
SEQ ID No. 490: 5'- GTTATCCCCTGCTAAGAG  
SEQ ID No. 491: 5'- CCCGTTCGCCACTCTTTG  
SEQ ID No. 492: 5'- AGCTGAGGGCTTTCACTT  
20 SEQ ID No. 493: 5'- GAGCTGAGGGCTTTCACT  
SEQ ID No. 494: 5'- GCTGAGGGCTTTCACTTC  
SEQ ID No. 495: 5'- CTGAGGGCTTTCACTTCA

The sequences SEQ ID No. 445 to SEQ ID No. 495 are particularly suitable for the  
25 detection of bacteria of the genus *Weissella*.

SEQ ID No. 496: 5' CCCGTGTCCCGAAGGAAC  
SEQ ID No. 497: 5' GCACGAGTATGTCAAGAC  
SEQ ID No. 498: 5' GTATCCCGTGTCCCGAAG  
30 SEQ ID No. 499: 5' TCCCGTGTCCCGAAGGAA

SEQ ID No. 500: 5' ATCCCGTGTCCCGAAGGA  
SEQ ID No. 501: 5' TATCCCGTGTCCCGAAGG  
SEQ ID No. 502: 5' CTTACCTTAGGAAGCGCC  
SEQ ID No. 503: 5' TTACCTTAGGAAGCGCCC  
5 SEQ ID No. 504: 5' CCTGTATCCCGTGTCCCG  
SEQ ID No. 505: 5' CCACCTGTATCCCGTGTC  
SEQ ID No. 506: 5' CACCTGTATCCCGTGTCC  
SEQ ID No. 507: 5' ACCTGTATCCCGTGTCCC  
SEQ ID No. 508: 5' CTGTATCCCGTGTCCCGA  
10 SEQ ID No. 509: 5' TGTATCCCGTGTCCCGAA  
SEQ ID No. 510: 5' CACGAGTATGTCAAGACC  
SEQ ID No. 511: 5' CGGTCTTACCTTAGGAAG  
SEQ ID No. 512: 5' TAGGAAGCGCCCTCCTTG  
SEQ ID No. 513: 5' AGGAAGCGCCCTCCTTGC  
15 SEQ ID No. 514: 5' TTAGGAAGCGCCCTCCTT  
SEQ ID No. 515: 5' CTTAGGAAGCGCCCTCCT  
SEQ ID No. 516: 5' CCTTAGGAAGCGCCCTCC  
SEQ ID No. 517: 5' ACCTTAGGAAGCGCCCTC  
SEQ ID No. 518: 5' TGCACACAATGGTTGAGC  
20 SEQ ID No. 519: 5' TACCTTAGGAAGCGCCCT  
SEQ ID No. 520: 5' ACCACCTGTATCCCGTGT  
SEQ ID No. 521: 5' GCACCACCTGTATCCCGT  
SEQ ID No. 522: 5' CACCACCTGTATCCCGTG  
SEQ ID No. 523: 5' GCGGTTAGGCAACCTACT  
25 SEQ ID No. 524: 5' TGCGGTTAGGCAACCTAC  
SEQ ID No. 525: 5' TTGCGGTTAGGCAACCTA  
SEQ ID No. 526: 5' GGTCTTACCTTAGGAAGC  
SEQ ID No. 527: 5' GCTAATACAACGCGGGAT  
SEQ ID No. 528: 5' CTAATACAACGCGGGATC  
30 SEQ ID No. 529: 5' ATACAACGCGGGATCATC

SEQ ID No. 530: 5' CGGTTAGGCAACCTACTT  
SEQ ID No. 531: 5' TGCACCACCTGTATCCCG  
SEQ ID No. 532: 5' GAAGCGCCCTCCTTGCGG  
SEQ ID No. 533: 5' GGAAGCGCCCTCCTTGCG  
5 SEQ ID No. 534: 5' CGTCCCTTTCTGGTTAGA  
SEQ ID No. 535: 5' AGCTAATACAACGCGGGA  
SEQ ID No. 536: 5' TAGCTAATACAACGCGGG  
SEQ ID No. 537: 5' CTAGCTAATACAACGCGG  
SEQ ID No. 538: 5' GGCTATGTATCATCGCCT  
10 SEQ ID No. 539: 5' GAGCCACTGCCTTTTACA  
SEQ ID No. 540: 5' GTCGGCTATGTATCATCG  
SEQ ID No. 541: 5' GGTCGGCTATGTATCATC  
SEQ ID No. 542: 5' CAGGTCGGCTATGTATCA  
SEQ ID No. 543: 5' CGGCTATGTATCATCGCC  
15 SEQ ID No. 544: 5' TCGGCTATGTATCATCGC  
SEQ ID No. 545: 5' GTCTTACCTTAGGAAGCG  
SEQ ID No. 546: 5' TCTTACCTTAGGAAGCGC

The sequences SEQ ID No. 496 to SEQ ID No. 546 are particularly suitable for the  
20 detection of bacteria of the genus *Lactococcus*.

d) Nucleic acid molecules, which specifically detect drink-spoiling acetic acid  
bacteria:

25 SEQ ID No. 547: 5'- GTACAAACCGCCTACACGCC  
SEQ ID No. 548: 5'- TGTACAAACCGCCTACACGC  
SEQ ID No. 549: 5'- GATCAGCACGATGTCGCCAT  
SEQ ID No. 550: 5'- CTGTACAAACCGCCTACACG  
SEQ ID No. 551: 5'- GAGATCAGCACGATGTCGCC  
30 SEQ ID No. 552: 5'- AGATCAGCACGATGTCGCCA

SEQ ID No. 553: 5'- ATCAGCACGATGTCGCCATC  
SEQ ID No. 554: 5'- TCAGCACGATGTCGCCATCT  
SEQ ID No. 555: 5'- ACTGTACAAACCGCCTACAC  
SEQ ID No. 556: 5'- CCGCCACTAAGGCCGAAACC  
5 SEQ ID No. 557: 5'- CAGCACGATGTCGCCATCTA  
SEQ ID No. 558: 5'- TACAAACCGCCTACACGCCC  
SEQ ID No. 559: 5'- AGCACGATGTCGCCATCTAG  
SEQ ID No. 560: 5'- CGGCTTTTAGAGATCAGCAC  
SEQ ID No. 561: 5'- TCCGCCACTAAGGCCGAAAC  
10 SEQ ID No. 562: 5'- GACTGTACAAACCGCCTACA  
SEQ ID No. 563: 5'- GTCCGCCACTAAGGCCGAAA  
SEQ ID No. 564: 5'- GGGGATTTACATCTGACTG  
SEQ ID No. 565: 5'- CATACAAGCCCTGGTAAGGT  
SEQ ID No. 566: 5'- ACAAGCCCTGGTAAGGTTCT  
15 SEQ ID No. 567: 5'- ACAAACCGCCTACACGCCCT  
SEQ ID No. 568: 5'- CTGACTGTACAAACCGCCTA  
SEQ ID No. 569: 5'- TGACTGTACAAACCGCCTAC  
SEQ ID No. 570: 5'- ACGATGTCGCCATCTAGCTT  
SEQ ID No. 571: 5'- CACGATGTCGCCATCTAGCT  
20 SEQ ID No. 572: 5'- CGATGTCGCCATCTAGCTTC  
SEQ ID No. 573: 5'- GCACGATGTCGCCATCTAGC  
SEQ ID No. 574: 5'- GATGTCGCCATCTAGCTTCC  
SEQ ID No. 575: 5'- ATGTCGCCATCTAGCTTCCC  
SEQ ID No. 576: 5'- TGTCGCCATCTAGCTTCCCA  
25 SEQ ID No. 577: 5'- GCCATCTAGCTTCCCCTGT  
SEQ ID No. 578: 5'- TCGCCATCTAGCTTCCCCT  
SEQ ID No. 579: 5'- CGCCATCTAGCTTCCCCTG  
SEQ ID No. 580: 5'- GTCGCCATCTAGCTTCCCAC  
SEQ ID No. 581: 5'- TACAAGCCCTGGTAAGGTTT  
30 SEQ ID No. 582: 5'- GCCACTAAGGCCGAAACCTT

	SEQ ID No. 583:	5'- ACTAAGGCCGAAACCTTCGT
	SEQ ID No. 584:	5'- CTAAGGCCGAAACCTTCGTG
	SEQ ID No. 585:	5'- CACTAAGGCCGAAACCTTCG
	SEQ ID No. 586:	5'- AAGGCCGAAACCTTCGTGCG
5	SEQ ID No. 587:	5'- CCACTAAGGCCGAAACCTTC
	SEQ ID No. 588:	5'- TAAGGCCGAAACCTTCGTGC
	SEQ ID No. 589:	5'- AGGCCGAAACCTTCGTGCGA
	SEQ ID No. 590:	5'- TCTGACTGTACAAACCGCCT
	SEQ ID No. 591:	5'- CATCTGACTGTACAAACCGC
10	SEQ ID No. 592:	5'- ATCTGACTGTACAAACCGCC
	SEQ ID No. 593:	5'- CTTCGTGCGACTTG CATGTG
	SEQ ID No. 594:	5'- CCTTCGTGCGACTTG CATGT
	SEQ ID No. 595:	5'- CTCTCTAGAGTGCCCAACCA
	SEQ ID No. 596:	5'- TCTCTAGAGTGCCCAACCAA
15	SEQ ID No. 597:	5'- ACGTATCAAATGCAGCTCCC
	SEQ ID No. 598:	5'- CGTATCAAATGCAGCTCCCA
	SEQ ID No. 599:	5'- CGCCACTAAGGCCGAAACCT
	SEQ ID No. 600:	5'- CCGAAACCTTCGTGCGACTT
	SEQ ID No. 601:	5'- GCCGAAACCTTCGTGCGACT
20	SEQ ID No. 602:	5'- AACCTTCGTGCGACTTG CAT
	SEQ ID No. 603:	5'- CGAAACCTTCGTGCGACTTG
	SEQ ID No. 604:	5'- ACCTTCGTGCGACTTG CATG
	SEQ ID No. 605:	5'- GAAACCTTCGTGCGACTTG C
	SEQ ID No. 606:	5'- GGCCGAAACCTTCGTGCGAC
25	SEQ ID No. 607:	5'- AAACCTTCGTGCGACTTGCA
	SEQ ID No. 608:	5'- CACGTATCAAATGCAGCTCC

The sequences SEQ ID No. 547 to SEQ ID No. 608 are particularly suitable for the simultaneous detection of bacteria of the genera *Acetobacter* and *Gluconobacter*.

SEQ ID No. 609: 5'- GCTCACCGGCTTAAGGTCAA  
SEQ ID No. 610: 5'- CGCTCACCGGCTTAAGGTCA  
SEQ ID No. 611: 5'- TCGCTCACCGGCTTAAGGTC  
SEQ ID No. 612: 5'- CTCACCGGCTTAAGGTCAAA  
5 SEQ ID No. 613: 5'- CCCGACCGTGGTCGGCTGCG  
SEQ ID No. 614: 5'- GCTCACCGGCTTAAGGTCAA  
SEQ ID No. 615: 5'- CGCTCACCGGCTTAAGGTCA  
SEQ ID No. 616: 5'- TCGCTCACCGGCTTAAGGTC  
SEQ ID No. 617: 5'- CTCACCGGCTTAAGGTCAAA  
10 SEQ ID No. 618: 5'- CCCGACCGTGGTCGGCTGCG  
SEQ ID No. 619: 5'- TCACCGGCTTAAGGTCAAAC  
SEQ ID No. 620: 5'- CAACCCTCTCTCACACTCTA  
SEQ ID No. 621: 5'- ACAACCCTCTCTCACACTCT  
SEQ ID No. 622: 5'- CCACAACCCTCTCTCACACT  
15 SEQ ID No. 623: 5'- AACCCTCTCTCACACTCTAG  
SEQ ID No. 624: 5'- CACAACCCTCTCTCACACTC  
SEQ ID No. 625: 5'- TCCACAACCCTCTCTCACAC  
SEQ ID No. 626: 5'- TTCCACAACCCTCTCTCACA  
SEQ ID No. 627: 5'- ACCCTCTCTCACACTCTAGT  
20 SEQ ID No. 628: 5'- GAGCCAGGTTGCCGCCTTCG  
SEQ ID No. 629: 5'- AGGTCAAACCAACTCCCATG  
SEQ ID No. 630: 5'- ATGAGCCAGGTTGCCGCCTT  
SEQ ID No. 631: 5'- TGAGCCAGGTTGCCGCCTTC  
SEQ ID No. 632: 5'- AGGCTCCTCCACAGGCGACT  
25 SEQ ID No. 633: 5'- CAGGCTCCTCCACAGGCGAC  
SEQ ID No. 634: 5'- GCAGGCTCCTCCACAGGCGA  
SEQ ID No. 635: 5'- TTCGCTCACCGGCTTAAGGT  
SEQ ID No. 636: 5'- GTTCGCTCACCGGCTTAAGG  
SEQ ID No. 637: 5'- GGTTGCTCACCGGCTTAAG  
30 SEQ ID No. 638: 5'- ATTCCACAACCCTCTCTCAC



	SEQ ID No. 639:	5'- TGACCCGACCGTGGTCGGCT
	SEQ ID No. 640:	5'- CCCTCTCTCACACTCTAGTC
	SEQ ID No. 641:	5'- GAATTCCACAACCCTCTCTC
	SEQ ID No. 642:	5'- AGCCAGGTTGCCGCCTTCGC
5	SEQ ID No. 643:	5'- GCCAGGTTGCCGCCTTCGCC
	SEQ ID No. 644:	5'- GGAATTCCACAACCCTCTCT
	SEQ ID No. 645:	5'- GGGAATTCCACAACCCTCTC
	SEQ ID No. 646:	5'- AACGCAGGCTCCTCCACAGG
	SEQ ID No. 647:	5'- CGGCTTAAGGTCAAACCAAC
10	SEQ ID No. 648:	5'- CCGGCTTAAGGTCAAACCAA
	SEQ ID No. 649:	5'- CACCGGCTTAAGGTCAAACC
	SEQ ID No. 650:	5'- ACCGGCTTAAGGTCAAACCA
	SEQ ID No. 651:	5'- ACCCAACATCCAGCACACAT
	SEQ ID No. 652:	5'- TCGCTGACCCGACCGTGGTC
15	SEQ ID No. 653:	5'- CGCTGACCCGACCGTGGTCG
	SEQ ID No. 654:	5'- GACCCGACCGTGGTCGGCTG
	SEQ ID No. 655:	5'- GCTGACCCGACCGTGGTCGG
	SEQ ID No. 656:	5'- CTGACCCGACCGTGGTCGGC
	SEQ ID No. 657:	5'- CAGGCGACTTGCGCCTTTGA
20	SEQ ID No. 658:	5'- TCATGCGGTATTAGCTCCAG
	SEQ ID No. 659:	5'- ACTAGCTAATCGAACGCAGG
	SEQ ID No. 660:	5'- CATGCGGTATTAGCTCCAGT
	SEQ ID No. 661:	5'- CGCAGGCTCCTCCACAGGCG
	SEQ ID No. 662:	5'- ACGCAGGCTCCTCCACAGGC
25	SEQ ID No. 663:	5'- CTCAGGTGTCATGCGGTATT
	SEQ ID No. 664:	5'- CGCCTTTGACCCTCAGGTGT
	SEQ ID No. 665:	5'- ACCCTCAGGTGTCATGCGGT
	SEQ ID No. 666:	5'- CCTCAGGTGTCATGCGGTAT
	SEQ ID No. 667:	5'- TTTGACCCTCAGGTGTCATG
30	SEQ ID No. 668:	5'- GACCCTCAGGTGTCATGCGG

	SEQ ID No. 669:	5'- TGACCCTCAGGTGTCATGCG
	SEQ ID No. 670:	5'- GCCTTTGACCCTCAGGTGTC
	SEQ ID No. 671:	5'- TTGACCCTCAGGTGTCATGC
	SEQ ID No. 672:	5'- CCCTCAGGTGTCATGCGGTA
5	SEQ ID No. 673:	5'- CCTTTGACCCTCAGGTGTCA
	SEQ ID No. 674:	5'- CTTTGACCCTCAGGTGTCAT
	SEQ ID No. 675:	5'- AGTTATCCCCCACCCTATGGA
	SEQ ID No. 676:	5'- CCAGCTATCGATCATCGCCT
	SEQ ID No. 677:	5'- ACCAGCTATCGATCATCGCC
10	SEQ ID No. 678:	5'- CAGCTATCGATCATCGCCTT
	SEQ ID No. 679:	5'- AGCTATCGATCATCGCCTTG
	SEQ ID No. 680:	5'- GCTATCGATCATCGCCTTGG
	SEQ ID No. 681:	5'- CTATCGATCATCGCCTTGGT
	SEQ ID No. 682:	5'- TTCGTGCGACTTGTCATGTGT
15	SEQ ID No. 683:	5'- TCGATCATCGCCTTGGTAGG
	SEQ ID No. 684:	5'- ATCGATCATCGCCTTGGTAG
	SEQ ID No. 685:	5'- CACAGGCGACTTGCGCCTTT
	SEQ ID No. 686:	5'- CCACAGGCGACTTGCGCCTT
	SEQ ID No. 687:	5'- TCCACAGGCGACTTGCGCCT
20	SEQ ID No. 688:	5'- TCCTCCACAGGCGACTTGCG
	SEQ ID No. 689:	5'- CCTCCACAGGCGACTTGCGC
	SEQ ID No. 690:	5'- CTCCACAGGCGACTTGCGCC
	SEQ ID No. 691:	5'- ACAGGCGACTTGCGCCTTTG
	SEQ ID No. 692:	5'- GCTCACCGGCTTAAGGTCAA
25	SEQ ID No. 693:	5'- CGCTCACCGGCTTAAGGTCA
	SEQ ID No. 694:	5'- TCGCTCACCGGCTTAAGGTC
	SEQ ID No. 695:	5'- CTCACCGGCTTAAGGTCAAA
	SEQ ID No. 696:	5'- CCCGACCGTGGTCGGCTGCG
	SEQ ID No. 697:	5'- TCACCGGCTTAAGGTCAAAC
30	SEQ ID No. 698:	5'- CAACCCTCTCTCACACTCTA

SEQ ID No. 699: 5'- ACAACCCTCTCTCACACTCT  
SEQ ID No. 700: 5'- CCACAACCCTCTCTCACACT  
SEQ ID No. 701: 5'- AACCTCTCTCTCACACTCTAG  
SEQ ID No. 702: 5'- CACAACCCTCTCTCACACTC  
5 SEQ ID No. 703: 5'- TCCACAACCCTCTCTCACAC  
SEQ ID No. 704: 5'- TTCCACAACCCTCTCTCACA  
SEQ ID No. 705: 5'- ACCCTCTCTCACACTCTAGT  
SEQ ID No. 706: 5'- GAGCCAGGTTGCCGCCTTCG  
SEQ ID No. 707: 5'- AGGTCAAACCAACTCCCATG  
10 SEQ ID No. 708: 5'- ATGAGCCAGGTTGCCGCCTT  
SEQ ID No. 709: 5'- TGAGCCAGGTTGCCGCCTTC  
SEQ ID No. 710: 5'- AGGCTCCTCCACAGGCGACT  
SEQ ID No. 711: 5'- CAGGCTCCTCCACAGGCGAC  
SEQ ID No. 712: 5'- GCAGGCTCCTCCACAGGCGA  
15 SEQ ID No. 713: 5'- TTCGCTCACCGGCTTAAGGT  
SEQ ID No. 714: 5'- GTTCGCTCACCGGCTTAAGG  
SEQ ID No. 715: 5'- GGTTCGCTCACCGGCTTAAG  
SEQ ID No. 716: 5'- ATTCCACAACCCTCTCTCAC  
SEQ ID No. 717: 5'- TGACCCGACCGTGGTCGGCT  
20 SEQ ID No. 718: 5'- CCCTCTCTCACACTCTAGTC  
SEQ ID No. 719: 5'- GAATTCCACAACCCTCTCTC  
SEQ ID No. 720: 5'- AGCCAGGTTGCCGCCTTCGC  
SEQ ID No. 721: 5'- GCCAGGTTGCCGCCTTCGCC  
SEQ ID No. 722: 5'- GGAATTCCACAACCCTCTCT  
25 SEQ ID No. 723: 5'- GGGAATTCCACAACCCTCTC  
SEQ ID No. 724: 5'- AACGCAGGCTCCTCCACAGG  
SEQ ID No. 725: 5'- CGGCTTAAGGTCAAACCAAC  
SEQ ID No. 726: 5'- CCGGCTTAAGGTCAAACCAA  
SEQ ID No. 727: 5'- CACCGGCTTAAGGTCAAACC  
30 SEQ ID No. 728: 5'- ACCGGCTTAAGGTCAAACCA

	SEQ ID No. 729:	5'- ACCCAACATCCAGCACACAT
	SEQ ID No. 730:	5'- TCGCTGACCCGACCGTGGTC
	SEQ ID No. 731:	5'- CGCTGACCCGACCGTGGTCG
	SEQ ID No. 732:	5'- GACCCGACCGTGGTCGGCTG
5	SEQ ID No. 733:	5'- GCTGACCCGACCGTGGTCGG
	SEQ ID No. 734:	5'- CTGACCCGACCGTGGTCGGC
	SEQ ID No. 735:	5'- CAGGCGACTTGCGCCTTTGA
	SEQ ID No. 736:	5'- TCATGCGGTATTAGCTCCAG
	SEQ ID No. 737:	5'- ACTAGCTAATCGAACGCAGG
10	SEQ ID No. 738:	5'- CATGCGGTATTAGCTCCAGT
	SEQ ID No. 739:	5'- CGCAGGCTCCTCCACAGGCG
	SEQ ID No. 740:	5'- ACGCAGGCTCCTCCACAGGC
	SEQ ID No. 741:	5'- CTCAGGTGTCATGCGGTATT
	SEQ ID No. 742:	5'- CGCCTTTGACCCTCAGGTGT
15	SEQ ID No. 743:	5'- ACCCTCAGGTGTCATGCGGT
	SEQ ID No. 744:	5'- CCTCAGGTGTCATGCGGTAT
	SEQ ID No. 745:	5'- TTTGACCCTCAGGTGTCATG
	SEQ ID No. 746:	5'- GACCCTCAGGTGTCATGCGG
	SEQ ID No. 747:	5'- TGACCCTCAGGTGTCATGCG
20	SEQ ID No. 748:	5'- GCCTTTGACCCTCAGGTGTC
	SEQ ID No. 749:	5'- TTGACCCTCAGGTGTCATGC
	SEQ ID No. 750:	5'- CCCTCAGGTGTCATGCGGTA
	SEQ ID No. 751:	5'- CCTTTGACCCTCAGGTGTCA
	SEQ ID No. 752:	5'- CTTTGACCCTCAGGTGTCAT
25	SEQ ID No. 753:	5'- AGTTATCCCCCACCATGGA
	SEQ ID No. 754:	5'- CCAGCTATCGATCATCGCCT
	SEQ ID No. 755:	5'- ACCAGCTATCGATCATCGCC
	SEQ ID No. 756:	5'- CAGCTATCGATCATCGCCTT
	SEQ ID No. 757:	5'- AGCTATCGATCATCGCCTTG
30	SEQ ID No. 758:	5'- GCTATCGATCATCGCCTTGG

SEQ ID No. 759: 5'- CTATCGATCATCGCCTTGGT  
SEQ ID No. 760: 5'- TTCGTGCGACTTGCGCATGTGT  
SEQ ID No. 761: 5'- TCGATCATCGCCTTGGTAGG  
SEQ ID No. 762: 5'- ATCGATCATCGCCTTGGTAG  
5 SEQ ID No. 763: 5'- CACAGGCGACTTGCGCCTTT  
SEQ ID No. 764: 5'- CCACAGGCGACTTGCGCCTT  
SEQ ID No. 765: 5'- TCCACAGGCGACTTGCGCCT  
SEQ ID No. 766: 5'- TCCTCCACAGGCGACTTGCG  
SEQ ID No. 767: 5'- CCTCCACAGGCGACTTGCGC  
10 SEQ ID No. 768: 5'- CTCCACAGGCGACTTGCGCC  
SEQ ID No. 769: 5'- ACAGGCGACTTGCGCCTTTG  
SEQ ID No. 770: 5'- TCACCGGCTTAAGGTCAAAC  
SEQ ID No. 771: 5'- CAACCCTCTCTCACACTCTA  
SEQ ID No. 772: 5'- ACAACCCTCTCTCACACTCT  
15 SEQ ID No. 773: 5'- CCACAACCCTCTCTCACACT  
SEQ ID No. 774: 5'- AACCCTCTCTCACACTCTAG  
SEQ ID No. 775: 5'- CACAACCCTCTCTCACACTC  
SEQ ID No. 776: 5'- TCCACAACCCTCTCTCACAC  
SEQ ID No. 777: 5'- TTCCACAACCCTCTCTCACA  
20 SEQ ID No. 778: 5'- ACCCTCTCTCACACTCTAGT  
SEQ ID No. 779: 5'- GAGCCAGGTTGCCGCCTTCG  
SEQ ID No. 780: 5'- AGGTCAAACCAACTCCCATG  
SEQ ID No. 781: 5'- ATGAGCCAGGTTGCCGCCTT  
SEQ ID No. 782: 5'- TGAGCCAGGTTGCCGCCTTC  
25 SEQ ID No. 783: 5'- AGGCTCCTCCACAGGCGACT  
SEQ ID No. 784: 5'- CAGGCTCCTCCACAGGCGAC  
SEQ ID No. 785: 5'- GCAGGCTCCTCCACAGGCGA  
SEQ ID No. 786: 5'- TTCGCTCACCGGCTTAAGGT  
SEQ ID No. 787: 5'- GTTCGCTCACCGGCTTAAGG  
30 SEQ ID No. 788: 5'- GGTTGCTCACCGGCTTAAG

SEQ ID No. 789: 5'- ATTCCACAACCCTCTCTCAC  
SEQ ID No. 790: 5'- TGACCCGACCGTGGTCGGCT  
SEQ ID No. 791: 5'- CCCTCTCTCACACTCTAGTC  
SEQ ID No. 792: 5'- GAATTCCACAACCCTCTCTC  
5 SEQ ID No. 793: 5'- AGCCAGGTTGCCGCCTTCGC  
SEQ ID No. 794: 5'- GCCAGGTTGCCGCCTTCGCC  
SEQ ID No. 795: 5'- GGAATTCCACAACCCTCTCT  
SEQ ID No. 796: 5'- GGGAATTCCACAACCCTCTC  
SEQ ID No. 797: 5'- AACGCAGGCTCCTCCACAGG  
10 SEQ ID No. 798: 5'- CGGCTTAAGGTCAAACCAAC  
SEQ ID No. 799: 5'- CCGGCTTAAGGTCAAACCAA  
SEQ ID No. 800: 5'- CACCGGCTTAAGGTCAAACC  
SEQ ID No. 801: 5'- ACCGGCTTAAGGTCAAACCA  
SEQ ID No. 802: 5'- ACCCAACATCCAGCACACAT  
15 SEQ ID No. 803: 5'- TCGCTGACCCGACCGTGGTC  
SEQ ID No. 804: 5'- CGCTGACCCGACCGTGGTCG  
SEQ ID No. 805: 5'- GACCCGACCGTGGTCGGCTG  
SEQ ID No. 806: 5'- GCTGACCCGACCGTGGTCGG  
SEQ ID No. 807: 5'- CTGACCCGACCGTGGTCGGC  
20 SEQ ID No. 808: 5'- CAGGCGACTTGCGCCTTTGA  
SEQ ID No. 809: 5'- TCATGCGGTATTAGCTCCAG  
SEQ ID No. 810: 5'- ACTAGCTAATCGAACGCAGG  
SEQ ID No. 811: 5'- CATGCGGTATTAGCTCCAGT  
SEQ ID No. 812: 5'- CGCAGGCTCCTCCACAGGCG  
25 SEQ ID No. 813: 5'- ACGCAGGCTCCTCCACAGGC  
SEQ ID No. 814: 5'- CTCAGGTGTCATGCGGTATT  
SEQ ID No. 815: 5'- CGCCTTTGACCCTCAGGTGT  
SEQ ID No. 816: 5'- ACCCTCAGGTGTCATGCGGT  
SEQ ID No. 817: 5'- CCTCAGGTGTCATGCGGTAT  
30 SEQ ID No. 818: 5'- TTTGACCCTCAGGTGTCATG

	SEQ ID No. 819:	5'- GACCCTCAGGTGTCATGCGG
	SEQ ID No. 820:	5'- TGACCCTCAGGTGTCATGCG
	SEQ ID No. 821:	5'- GCCTTTGACCCTCAGGTGTC
	SEQ ID No. 822:	5'- TTGACCCTCAGGTGTCATGC
5	SEQ ID No. 823:	5'- CCCTCAGGTGTCATGCGGTA
	SEQ ID No. 824:	5'- CCTTTGACCCTCAGGTGTCA
	SEQ ID No. 825:	5'- CTTTGACCCTCAGGTGTCAT
	SEQ ID No. 826:	5'- AGTTATCCCCCACCATGGA
	SEQ ID No. 827:	5'- CCAGCTATCGATCATCGCCT
10	SEQ ID No. 828:	5'- ACCAGCTATCGATCATCGCC
	SEQ ID No. 829:	5'- CAGCTATCGATCATCGCCTT
	SEQ ID No. 830:	5'- AGCTATCGATCATCGCCTTG
	SEQ ID No. 831:	5'- GCTATCGATCATCGCCTTGG
	SEQ ID No. 832:	5'- CTATCGATCATCGCCTTGGT
15	SEQ ID No. 833:	5'- TTCGTGCGACTTGTCATGTGT
	SEQ ID No. 834:	5'- TCGATCATCGCCTTGGTAGG
	SEQ ID No. 835:	5'- ATCGATCATCGCCTTGGTAG
	SEQ ID No. 836:	5'- CACAGGCGACTTGCGCCTTT
	SEQ ID No. 837:	5'- CCACAGGCGACTTGCGCCTT
20	SEQ ID No. 838:	5'- TCCACAGGCGACTTGCGCCT
	SEQ ID No. 839:	5'- TCCTCCACAGGCGACTTGCG
	SEQ ID No. 840:	5'- CCTCCACAGGCGACTTGCGC
	SEQ ID No. 841:	5'- CTCCACAGGCGACTTGCGCC
	SEQ ID No. 842:	5'- ACAGGCGACTTGCGCCTTTG

25

The sequences SEQ ID No. 609 to SEQ ID No. 842 are particularly suitable for the simultaneous detection of bacteria of the genera *Acetobacter*, *Gluconobacter* and *Gluconoacetobacter*.

30 e) Nucléic acid probe molecules, which specifically detect drink-spoiling bacilli:

SEQ ID No. 843: 5'- AGCCCCGGTTTCCCGGCGTT  
SEQ ID No. 844: 5'- CGCCTTTCCTTTTTCCTCCA  
SEQ ID No. 845: 5'- GCCCCGGTTTCCCGGCGTTA  
5 SEQ ID No. 846: 5'- GCCGCCTTTCCTTTTTCCTC  
SEQ ID No. 847: 5'- TAGCCCCGGTTTCCCGGCGT  
SEQ ID No. 848: 5'- CCGGGTACCGTCAAGGCGCC  
SEQ ID No. 849: 5'- AAGCCGCCTTTCCTTTTTC  
SEQ ID No. 850: 5'- CCCC GGTTTCCCGGCGTTAT  
10 SEQ ID No. 851: 5'- CCGGCGTTATCCCAGTCTTA  
SEQ ID No. 852: 5'- AGCCGCCTTTCCTTTTTCCT  
SEQ ID No. 853: 5'- CCGCCTTTCCTTTTTCCTCC  
SEQ ID No. 854: 5'- TTAGCCCCGGTTTCCCGGCG  
SEQ ID No. 855: 5'- CCCGGCGTTATCCCAGTCTT  
15 SEQ ID No. 856: 5'- GCCGGGTACCGTCAAGGCGC  
SEQ ID No. 857: 5'- GGCCGGGTACCGTCAAGGCG  
SEQ ID No. 858: 5'- TCCCGGCGTTATCCCAGTCT  
SEQ ID No. 859: 5'- TGGCCGGGTACCGTCAAGGC  
SEQ ID No. 860: 5'- GAAGCCGCCTTTCCTTTTTC  
20 SEQ ID No. 861: 5'- CCCGGTTTCCCGGCGTTATC  
SEQ ID No. 862: 5'- CGGCGTTATCCCAGTCTTAC  
SEQ ID No. 863: 5'- GGCGTTATCCCAGTCTTACA  
SEQ ID No. 864: 5'- GCGTTATCCCAGTCTTACAG  
SEQ ID No. 865: 5'- CGGGTACCGTCAAGGCGCCG  
25 SEQ ID No. 866: 5'- ATTAGCCCCGGTTTCCCGGC  
SEQ ID No. 867: 5'- AAGGGGAAGGCCCTGTCTCC  
SEQ ID No. 868: 5'- GGCCCTGTCTCCAGGGAGGT  
SEQ ID No. 869: 5'- AGGCCCTGTCTCCAGGGAGG  
SEQ ID No. 870: 5'- AAGGCCCTGTCTCCAGGGAG  
30 SEQ ID No. 871: 5'- GCCCTGTCTCCAGGGAGGTC



	SEQ ID No. 872:	5'- CGTTATCCCAGTCTTACAGG
	SEQ ID No. 873:	5'- GGGTACCGTCAAGGCGCCGC
	SEQ ID No. 874:	5'- CGGCAACAGAGTTTTACGAC
	SEQ ID No. 875:	5'- GGGGAAGGCCCTGTCTCCAG
5	SEQ ID No. 876:	5'- AGGGGAAGGCCCTGTCTCCA
	SEQ ID No. 877:	5'- GCAGCCGAAGCCGCCTTTCC
	SEQ ID No. 878:	5'- TTCTTCCCCGGCAACAGAGT
	SEQ ID No. 879:	5'- CGGCACTTGTTCTTCCCCGG
	SEQ ID No. 880:	5'- GTTCTTCCCCGGCAACAGAG
10	SEQ ID No. 881:	5'- GGC ACTTGTTCTTCCCCGGC
	SEQ ID No. 882:	5'- GCACTTGTTCTTCCCCGGCA
	SEQ ID No. 883:	5'- CACTTGTTCTTCCCCGGCAA
	SEQ ID No. 884:	5'- TCTTCCCCGGCAACAGAGTT
	SEQ ID No. 885:	5'- TTGTTCTTCCCCGGCAACAG
15	SEQ ID No. 886:	5'- ACTTGTTCTTCCCCGGCAAC
	SEQ ID No. 887:	5'- TGTTCTTCCCCGGCAACAGA
	SEQ ID No. 888:	5'- CTTGTTCTTCCCCGGCAACA
	SEQ ID No. 889:	5'- ACGGCACTTGTTCTTCCCCG
	SEQ ID No. 890:	5'- GTCCGCCGCTAACCTTTTAA
20	SEQ ID No. 891:	5'- CTGGCCGGGTACCGTCAAGG
	SEQ ID No. 892:	5'- TCTGGCCGGGTACCGTCAAG
	SEQ ID No. 893:	5'- TTCTGGCCGGGTACCGTCAA
	SEQ ID No. 894:	5'- CAATGCTGGCAACTAAGGTC
	SEQ ID No. 895:	5'- CGTCCGCCGCTAACCTTTTA
25	SEQ ID No. 896:	5'- CGAAGCCGCCTTTCCTTTTT
	SEQ ID No. 897:	5'- CCGAAGCCGCCTTTCCTTTT
	SEQ ID No. 898:	5'- GCCGAAGCCGCCTTTCCTTT
	SEQ ID No. 899:	5'- AGCCGAAGCCGCCTTTCCTT
	SEQ ID No. 900:	5'- ACCGTCAAGGCGCCGCCCTG
30	SEQ ID No. 901:	5'- CCGTGGCTTTCTGGCCGGGT

SEQ ID No. 902: 5'- GCTTTCTGGCCGGGTACCGT  
SEQ ID No. 903: 5'- GCCGTGGCTTTCTGGCCGGG  
SEQ ID No. 904: 5'- GGCTTTCTGGCCGGGTACCG  
SEQ ID No. 905: 5'- CTTTCTGGCCGGGTACCGTC  
5 SEQ ID No. 906: 5'- TGGCTTTCTGGCCGGGTACC  
SEQ ID No. 907: 5'- GTGGCTTTCTGGCCGGGTAC  
SEQ ID No. 908: 5'- CGTGGCTTTCTGGCCGGGTA  
SEQ ID No. 909: 5'- TTTCTGGCCGGGTACCGTCA  
SEQ ID No. 910: 5'- GGGAAGGCCCTGTCTCCAGG  
10 SEQ ID No. 911: 5'- CGAAGGGGAAGGCCCTGTCT  
SEQ ID No. 912: 5'- CCGAAGGGGAAGGCCCTGTC  
SEQ ID No. 913: 5'- GAAGGGGAAGGCCCTGTCTC  
SEQ ID No. 914: 5'- GGC GCCGCCCTGTTCGAACG  
SEQ ID No. 915: 5'- AGGCGCCGCCCTGTTCGAAC  
15 SEQ ID No. 916: 5'- AAGGCGCCGCCCTGTTCGAA  
SEQ ID No. 917: 5'- CCCGGCAACAGAGTTTTACG  
SEQ ID No. 918: 5'- CCCC GGCAACAGAGTTTTAC  
SEQ ID No. 919: 5'- CCATCTGTAAGTGGCAGCCG  
SEQ ID No. 920: 5'- TCTGTAAGTGGCAGCCGAAG  
20 SEQ ID No. 921: 5'- CTGTAAGTGGCAGCCGAAGC  
SEQ ID No. 922: 5'- C C C A T C T G T A A G T G G C A G C C  
SEQ ID No. 923: 5'- T G T A A G T G G C A G C C G A A G C C  
SEQ ID No. 924: 5'- C A T C T G T A A G T G G C A G C C G A  
SEQ ID No. 925: 5'- A T C T G T A A G T G G C A G C C G A A  
25 SEQ ID No. 926: 5'- C A G C C G A A G C C G C C T T T C C T  
SEQ ID No. 927: 5'- G G C A A C A G A G T T T T A C G A C C  
SEQ ID No. 928: 5'- C C G G C A A C A G A G T T T T A C G A  
SEQ ID No. 929: 5'- T T C C C C G G C A A C A G A G T T T T  
SEQ ID No. 930: 5'- C T T C C C C G G C A A C A G A G T T T  
30 SEQ ID No. 931: 5'- T C C C C G G C A A C A G A G T T T T A

SEQ ID No. 932: 5'- CCGTCCGCCGCTAACCTTTT

The sequences SEQ ID No. 843 to SEQ ID No. 932 are particularly suitable for the detection of *Bacillus coagulans*.

5

f) Nucleic acid probe molecules which specifically detect drink-spoiling Alicyclobacilli:

SEQ ID No. 933: 5'- CTCCTCCGACTTACGCCGG  
10 SEQ ID No. 934: 5'- CCTCCGACTTACGCCGGCAG  
SEQ ID No. 935: 5'- TTCCTCCGACTTACGCCGGC  
SEQ ID No. 936: 5'- TCCTCCGACTTACGCCGGCA  
SEQ ID No. 937: 5'- TCCGACTTACGCCGGCAGTC  
SEQ ID No. 938: 5'- CCGACTTACGCCGGCAGTCA  
15 SEQ ID No. 939: 5'- GCCTTCCTCCGACTTACGCC  
SEQ ID No. 940: 5'- CCTTCCTCCGACTTACGCCG  
SEQ ID No. 941: 5'- GCTCTCCCCGAGCAACAGAG  
SEQ ID No. 942: 5'- CTCTCCCCGAGCAACAGAGC  
SEQ ID No. 943: 5'- CGCTCTCCCCGAGCAACAGA  
20 SEQ ID No. 944: 5'- CTCCGACTTACGCCGGCAGT  
SEQ ID No. 945: 5'- TCTCCCCGAGCAACAGAGCT  
SEQ ID No. 946: 5'- CGACTTACGCCGGCAGTCAC  
SEQ ID No. 947: 5'- TCGGCACTGGGGTGTGTCCC  
SEQ ID No. 948: 5'- GGC ACTGGGGTGTGTCCCCC  
25 SEQ ID No. 949: 5'- CTGGGGTGTGTCCCCCAAC  
SEQ ID No. 950: 5'- CACTGGGGTGTGTCCCCCA  
SEQ ID No. 951: 5'- ACTGGGGTGTGTCCCCCAA  
SEQ ID No. 952: 5'- GCACTGGGGTGTGTCCCCC  
SEQ ID No. 953: 5'- TGGGGTGTGTCCCCCAACA  
30 SEQ ID No. 954: 5'- CACTCCAGACTTGCTCGACC

SEQ ID No. 955: 5'- TCACTCCAGACTTGCTCGAC  
SEQ ID No. 956: 5'- CGGCACTGGGGTGTGTCCCC  
SEQ ID No. 957: 5'- CGCCTTCCTCCGACTTACGC  
SEQ ID No. 958: 5'- CTCCCCGAGCAACAGAGCTT  
5 SEQ ID No. 959: 5'- ACTCCAGACTTGCTCGACCG  
SEQ ID No. 960: 5'- CCCATGCCGCTCTCCCCGAG  
SEQ ID No. 961: 5'- CCATGCCGCTCTCCCCGAGC  
SEQ ID No. 962: 5'- CCCCATGCCGCTCTCCCCGA  
SEQ ID No. 963: 5'- TCACTCGGTACCGTCTCGCA  
10 SEQ ID No. 964: 5'- CATGCCGCTCTCCCCGAGCA  
SEQ ID No. 965: 5'- ATGCCGCTCTCCCCGAGCAA  
SEQ ID No. 966: 5'- TTCGGCACTGGGGTGTGTCC  
SEQ ID No. 967: 5'- TGCCGCTCTCCCCGAGCAAC  
SEQ ID No. 968: 5'- TTCACTCCAGACTTGCTCGA  
15 SEQ ID No. 969: 5'- CCCGCAAGAAGATGCCTCCT  
SEQ ID No. 970: 5'- AGAAGATGCCTCCTCGCGGG  
SEQ ID No. 971: 5'- AAGAAGATGCCTCCTCGCGG  
SEQ ID No. 972: 5'- CGCAAGAAGATGCCTCCTCG  
SEQ ID No. 973: 5'- AAGATGCCTCCTCGCGGGCG  
20 SEQ ID No. 974: 5'- CCGCAAGAAGATGCCTCCTC  
SEQ ID No. 975: 5'- GAAAGATGCCTCCTCGCGGGC  
SEQ ID No. 976: 5'- CCCCAGCAAGAAGATGCCTCC  
SEQ ID No. 977: 5'- CAAGAAGATGCCTCCTCGCG  
SEQ ID No. 978: 5'- TCCTTCGGCACTGGGGTGTG  
25 SEQ ID No. 979: 5'- CCGCTCTCCCCGAGCAACAG  
SEQ ID No. 980: 5'- TGCCTCCTCGCGGGCGTATC  
SEQ ID No. 981: 5'- GACTTACGCCGGCAGTCACC  
SEQ ID No. 982: 5'- GGCTCCTCTCTCAGCGGCCC  
SEQ ID No. 983: 5'- CCTTCGGCACTGGGGTGTGT  
30 SEQ ID No. 984: 5'- GGGGTGTGTCCCCCAACAC

SEQ ID No. 985: 5'- GCCGCTCTCCCCGAGCAACA  
SEQ ID No. 986: 5'- AGATGCCTCCTCGCGGGCGT  
SEQ ID No. 987: 5'- CACTCGGTACCGTCTCGCAT  
SEQ ID No. 988: 5'- CTCACTCGGTACCGTCTCGC  
5 SEQ ID No. 989: 5'- GCAAGAAGATGCCTCCTCGC  
SEQ ID No. 990: 5'- CTCCAGACTTGCTCGACCGC  
SEQ ID No. 991: 5'- TTACGCCGGCAGTCACCTGT  
SEQ ID No. 992: 5'- CTTCGGCACTGGGGTGTGTC  
SEQ ID No. 993: 5'- CTCGCGGGCGTATCCGGCAT  
10 SEQ ID No. 994: 5'- GCCTCCTCGCGGGCGTATCC  
SEQ ID No. 995: 5'- ACTCGGTACCGTCTCGCATG  
SEQ ID No. 996: 5'- GATGCCTCCTCGCGGGCGTA  
SEQ ID No. 997: 5'- GGGTGTGTCCCCCAACACC  
SEQ ID No. 998: 5'- ACTTACGCCGGCAGTCACCT  
15 SEQ ID No. 999: 5'- CTTACGCCGGCAGTCACCTG  
SEQ ID No. 1000: 5'- ATGCCTCCTCGCGGGCGTAT  
SEQ ID No. 1001: 5'- GCGCCGCGGGCTCCTCTCTC  
SEQ ID No. 1002: 5'- GGTGTGTCCCCCAACACCT  
SEQ ID No. 1003: 5'- GTGTGTCCCCCAACACCTA  
20 SEQ ID No. 1004: 5'- CCTCGCGGGCGTATCCGGCA  
SEQ ID No. 1005: 5'- CCTCACTCGGTACCGTCTCG  
SEQ ID No. 1006: 5'- TCCTCACTCGGTACCGTCTC  
SEQ ID No. 1007: 5'- TCGCGGGCGTATCCGGCATT  
SEQ ID No. 1008: 5'- TTCACTCCAGACTTGCTCG  
25 SEQ ID No. 1009: 5'- TACGCCGGCAGTCACCTGTG  
SEQ ID No. 1010: 5'- TCCAGACTTGCTCGACCGCC  
SEQ ID No. 1011: 5'- CTCGGTACCGTCTCGCATGG  
SEQ ID No. 1012: 5'- CGCGGGCGTATCCGGCATTAA  
SEQ ID No. 1013: 5'- GCGTATCCGGCATTAGCGCC  
30 SEQ ID No. 1014: 5'- GGGCTCCTCTCTCAGCGGCC

SEQ ID No. 1015: 5'- TCCCCGAGCAACAGAGCTTT  
SEQ ID No. 1016: 5'- CCCCAGAGCAACAGAGCTTTA  
SEQ ID No. 1017: 5'- CCGAGCAACAGAGCTTTACA  
SEQ ID No. 1018: 5'- CCATCCCATGGTTGAGCCAT  
5 SEQ ID No. 1019: 5'- GTGTCCCCCAACACCTAGC  
SEQ ID No. 1020: 5'- GCGGGCGTATCCGGCATTAG  
SEQ ID No. 1021: 5'- CGAGCGGCTTTTTGGGTTC  
SEQ ID No. 1022: 5'- CTTTCACTCCAGACTTGCTC  
SEQ ID No. 1023: 5'- TTCCTTCGGCACTGGGGTGT  
10 SEQ ID No. 1024: 5'- CCGCCTTCCTCCGACTTACG  
SEQ ID No. 1025: 5'- CCCGCCTTCCTCCGACTTAC  
SEQ ID No. 1026: 5'- CCTCCTCGCGGGCGTATCCG  
SEQ ID No. 1027: 5'- TCCTCGCGGGCGTATCCGGC  
SEQ ID No. 1028: 5'- CATTAGCGCCCGTTTCCGGG  
15 SEQ ID No. 1029: 5'- GCATTAGCGCCCGTTTCCGG  
SEQ ID No. 1030: 5'- GGCATTAGCGCCCGTTTCCG  
SEQ ID No. 1031: 5'- GTCTCGCATGGGGCTTTCCA  
SEQ ID No. 1032: 5'- GCCATGGACTTTCACTCCAG  
SEQ ID No. 1033: 5'- CATGGACTTTCACTCCAGAC

20

The sequences SEQ ID No. 933 to SEQ ID No. 1033 are particularly suitable for the detection of bacteria of the genus *Alicyclobacillus*.

SEQ ID No. 1034: 5'- CCTTCCTCCGGCTTACGCCGGC  
25 SEQ ID No. 1035: 5'- CCTTCCTCCGACTTGCGCCGGC  
SEQ ID No. 1036: 5'- CCTTCCTCCGACTTTCACCGGC

The nucleic acid probe molecules according to SEQ ID No. 1034 to SEQ ID No. 1036 are used as unlabelled competitor probes for the detection of bacteria of the  
30 genus *Alicyclobacillus* in combination with the oligonucleotide probe according to

SEQ ID No. 933, in order to prevent the binding of the labelled oligonucleotide probe specific for bacteria of the genus *Alicyclobacillus* to nucleic acid sequences which are not specific for bacteria of the genus *Alicyclobacillus*.

- 5 SEQ ID No. 1037: 5'- ACCGTCTCACAAGGAGCTTT  
SEQ ID No. 1038: 5'- TACCGTCTCACAAGGAGCTT  
SEQ ID No. 1039: 5'- GTACCGTCTCACAAGGAGCT  
SEQ ID No. 1040: 5'- GCCTACCCGTGTATTATCCG  
SEQ ID No. 1041: 5'- CCGTCTCACAAGGAGCTTTC  
10 SEQ ID No. 1042: 5'- CTACCCGTGTATTATCCGGC  
SEQ ID No. 1043: 5'- GGTACCGTCTCACAAGGAGC  
SEQ ID No. 1044: 5'- CGTCTCACAAGGAGCTTTCC  
SEQ ID No. 1045: 5'- TCTCACAAGGAGCTTTCCAC  
SEQ ID No. 1046: 5'- TACCCGTGTATTATCCGGCA  
15 SEQ ID No. 1047: 5'- GTCTCACAAGGAGCTTTCCA  
SEQ ID No. 1048: 5'- ACCCGTGTATTATCCGGCAT  
SEQ ID No. 1049: 5'- CTCGGTACCGTCTCACAAGG  
SEQ ID No. 1050: 5'- CGGTACCGTCTCACAAGGAG  
SEQ ID No. 1051: 5'- ACTCGGTACCGTCTCACAAG  
20 SEQ ID No. 1052: 5'- CGGCTGGCTCCATAACGGTT  
SEQ ID No. 1053: 5'- ACAAGTAGATGECTACCCGT  
SEQ ID No. 1054: 5'- TGGCTECATAACGGTTACCT  
SEQ ID No. 1055: 5'- CAAGTAGATGCCTACCCGTG  
SEQ ID No. 1056: 5'- CACAAGTAGATGCCTACCCG  
25 SEQ ID No. 1057: 5'- GGCTCCATAACGGTTACCTC  
SEQ ID No. 1058: 5'- ACACAAGTAGATGCCTACCC  
SEQ ID No. 1059: 5'- CTGGCTCCATAACGGTTACC  
SEQ ID No. 1060: 5'- GCTGGCTCCATAACGGTTAC  
SEQ ID No. 1061: 5'- GGCTGGCTCCATAACGGTTA  
30 SEQ ID No. 1062: 5'- GCTCCATAACGGTTACCTCA

SEQ ID No. 1063: 5'- AAGTAGATGCCTACCCGTGT  
SEQ ID No. 1064: 5'- CTCCATAACGGTTACCTCAC  
SEQ ID No. 1065: 5'- TGCCTACCCGTGTATTATCC  
SEQ ID No. 1066: 5'- TCGGTACCGTCTCACAAGGA  
5 SEQ ID No. 1067: 5'- CTCACAAGGAGCTTTCCACT  
SEQ ID No. 1068: 5'- GTAGATGCCTACCCGTGTAT  
SEQ ID No. 1069: 5'- CCTACCCGTGTATTATCCGG  
SEQ ID No. 1070: 5'- CACTCGGTACCGTCTCACAA  
SEQ ID No. 1071: 5'- CTCAGCGATGCAGTTGCATC  
10 SEQ ID No. 1072: 5'- AGTAGATGCCTACCCGTGTA  
SEQ ID No. 1073: 5'- GCGGCTGGCTCCATAACGGT  
SEQ ID No. 1074: 5'- CCAAAGCAATCCCAAGGTTG  
SEQ ID No. 1075: 5'- TCCATAACGGTTACCTCACC  
SEQ ID No. 1076: 5'- CCCGTGTATTATCCGGCATT  
15 SEQ ID No. 1077: 5'- TCTCAGCGATGCAGTTGCAT  
SEQ ID No. 1078: 5'- CCATAACGGTTACCTCACCG  
SEQ ID No. 1079: 5'- TCAGCGATGCAGTTGCATCT  
SEQ ID No. 1080: 5'- GGCGGCTGGCTCCATAACGG  
SEQ ID No. 1081: 5'- AAGCAATCCCAAGGTTGAGC  
20 SEQ ID No. 1082: 5'- TCACTCGGTACCGTCTCACA  
SEQ ID No. 1083: 5'- CCGAGTGTTATTCCAGTCTG  
SEQ ID No. 1084: 5'- CACAAGGAGCTTTCCAETCT  
SEQ ID No. 1085: 5'- ACAAGGAGCTTTCCACTCTC  
SEQ ID No. 1086: 5'- TCACAAGGAGCTTTCCACTC  
25 SEQ ID No. 1087: 5'- CAGCGATGCAGTTGCATCTT  
SEQ ID No. 1088: 5'- CAAGGAGCTTTCCACTCTCC  
SEQ ID No. 1089: 5'- CCAGTCTGAAAGGCAGATTG  
SEQ ID No. 1090: 5'- CAGTCTGAAAGGCAGATTGC  
SEQ ID No. 1091: 5'- CGGCGGCTGGCTCCATAACG  
30 SEQ ID No. 1092: 5'- CCTCTCTCAGCGATGCAGTT



SEQ ID No. 1093: 5'- CTCTCTCAGCGATGCAGTTG  
SEQ ID No. 1094: 5'- TCTCTCAGCGATGCAGTTGC  
SEQ ID No. 1095: 5'- CTCTCAGCGATGCAGTTGCA  
SEQ ID No. 1096: 5'- CAATCCCAAGGTTGAGCCTT  
5 SEQ ID No. 1097: 5'- AATCCCAAGGTTGAGCCTTG  
SEQ ID No. 1098: 5'- AGCAATCCCAAGGTTGAGCC  
SEQ ID No. 1099: 5'- CTCACTCGGTACCGTCTCAC  
SEQ ID No. 1100: 5'- GCAATCCCAAGGTTGAGCCT  
SEQ ID No. 1101: 5'- GCCTTGGACTTTCACTTCAG  
10 SEQ ID No. 1102: 5'- CATAACGGTTACCTCACCGA  
SEQ ID No. 1103: 5'- CTCCTCTCTCAGCGATGCAG  
SEQ ID No. 1104: 5'- TCGGCGGCTGGCTCCATAAC  
SEQ ID No. 1105: 5'- AGTCTGAAAGGCAGATTGCC  
SEQ ID No. 1106: 5'- TCCTCTCTCAGCGATGCAGT  
15 SEQ ID No. 1107: 5'- CCCAAGGTTGAGCCTTGGAC  
SEQ ID No. 1108: 5'- ATAACGGTTACCTCACCGAC  
SEQ ID No. 1109: 5'- TCCCAAGGTTGAGCCTTGGA  
SEQ ID No. 1110: 5'- ATTATCCGGCATTAGCACCC  
SEQ ID No. 1111: 5'- CTACGTGCTGGTAACACAGA  
20 SEQ ID No. 1112: 5'- GCCGCTAGCCCCGAAGGGCT  
SEQ ID No. 1113: 5'- CTAGCCCCGAAGGGCTCGCT  
SEQ ID No. 1114: 5'- CGCTAGCCCCGAAGGGCTCG  
SEQ ID No. 1115: 5'- AGCCCCGAAGGGCTCGCTCG  
SEQ ID No. 1116: 5'- CCGCTAGCCCCGAAGGGCTC  
25 SEQ ID No. 1117: 5'- TAGCCCCGAAGGGCTCGCTC  
SEQ ID No. 1118: 5'- GCTAGCCCCGAAGGGCTCGC  
SEQ ID No. 1119: 5'- GCCCCGAAGGGCTCGCTCGA  
SEQ ID No. 1120: 5'- ATCCCAAGGTTGAGCCTTGG  
SEQ ID No. 1121: 5'- GAGCCTTGGACTTTCACTTC  
30 SEQ ID No. 1122: 5'- CAAGGTTGAGCCTTGGACTT

SEQ ID No. 1123: 5'- GAGCTTTCCACTCTCCTTGT  
SEQ ID No. 1124: 5'- CCAAGGTTGAGCCTTGGACT  
SEQ ID No. 1125: 5'- CGGGCTCCTCTCTCAGCGAT  
SEQ ID No. 1126: 5'- GGAGCTTTCCACTCTCCTTG  
5 SEQ ID No. 1127: 5'- GGGCTCCTCTCTCAGCGATG  
SEQ ID No. 1128: 5'- TCTCCTTGTCGCTCTCCCCG  
SEQ ID No. 1129: 5'- TCCTTGTCGCTCTCCCCGAG  
SEQ ID No. 1130: 5'- AGCTTTCCACTCTCCTTGTC  
SEQ ID No. 1131: 5'- CCACTCTCCTTGTCGCTCTC  
10 SEQ ID No. 1132: 5'- GGCTCCTCTCTCAGCGATGC  
SEQ ID No. 1133: 5'- CCTTGTCGCTCTCCCCGAGC  
SEQ ID No. 1134: 5'- CACTCTCCTTGTCGCTCTCC  
SEQ ID No. 1135: 5'- ACTCTCCTTGTCGCTCTCCC  
SEQ ID No. 1136: 5'- CTCTCCTTGTCGCTCTCCCC  
15 SEQ ID No. 1137: 5'- GCGGGCTCCTCTCTCAGCGA  
SEQ ID No. 1138: 5'- GGCTCCATCATGGTTACCTC

The sequences SEQ ID No. 1037 to SEQ ID No. 1138 are particularly suitable for the detection of *Alicyclobacillus acidoterrestris*.

20

SEQ ID No. 1139: 5'- CCGTCTCCTAAGGAGCTTTCCA...

The nucleic acid probe molecule according to SEQ ID No. 1139 is used as unlabelled competitor probe for the detection of *Alicyclobacillus acidoterrestris* in combination  
25 with the oligonucleotide probe according to SEQ ID No. 1044, in order to prevent the binding of the labelled oligonucleotide probe specific for *Alicyclobacillus acidoterrestris* to nucleic acid sequences which are not specific for *Alicyclobacillus acidoterrestris*.

30

SEQ ID No. 1140: 5'- TCCCTCCTTAACGGTTACCTCA...

SEQ ID No. 1141: 5'- TGGCTCCATAA(A/T)GGTTACCTCA

The nucleic acid probe molecules according to SEQ ID No. 1140 to SEQ ID No. 1141 are used as unlabelled competitor probe for the detection of *Alicyclobacillus acidoterrestris* in combination with the oligonucleotide probe according to SEQ ID No. 1057, in order to prevent the binding of the labelled oligonucleotide probe specific for *Alicyclobacillus acidoterrestris*, to nucleic acid sequences which are not specific for *Alicyclobacillus acidoterrestris*.

10 SEQ ID No. 1142: 5'- CTCCTCCGGCTTGCGCCGG

SEQ ID No. 1143: 5'- CGCTCTTCCCGA(G/T)TGACTGA

SEQ ID No. 1144: 5'- CCTCGGGCTCCTCCATC(A/T)GC

The sequences SEQ ID No. 1142 to SEQ ID No. 1144 are particularly suitable for the simultaneous detection of *Alicyclobacillus cycloheptanicus* and *A. herbarius*.

A further subject of the invention are derivatives of the above oligonucleotide sequences, demonstrating specific hybridization with target nucleic acid sequences of the respective microorganism despite deviations in sequence and/or length, and which are therefore suitable for use in a method according to the invention and ensure the specific detection of the respective microorganism. These derivatives especially include:

a) nucleic acid molecules which (i) are identical with respect to the bases to one of the above oligonucleotide sequences (SEQ ID No. 1, 5 to 146, 148 to 154, 157 to 160, 163 to 1033, 1037 to 1138, 1142 to 1144) to at least 80%, preferably to at least 90% particularly preferred to at least 92%, 94%, 96%, or (ii) differ from the above oligonucleotide sequences by one or more deletions and/or additions and which allow for a specific hybridization with nucleic acid sequences of drink-spoiling yeasts of the genera *Zygosaccharomyces*, *Hanseniaspora*, *Candida*,

*Brettanomyces*, *Dekkera*, *Pichia*, *Saccharomyces* and *Saccharomycodes* and in particular of the species *Zygosaccharomyces bailii*, *Z. mellis*, *Z. rouxii*, *Z. bisporus*, *Z. fermentati*, *Z. microellipsoides*, *Hanseniaspora uvarum*, *Candida intermedia*, *C. crusei* (*Issatchenkia orientalis*), *C. parapsilosis*, *Brettanomyces*  
5 *bruxellensis*, *B. naardenensis*, *Dekkera anomala*, *Pichia membranaefaciens*, *P. minuta*, *P. anomala*, *Saccharomyces exiguus*, *S. cerevisiae*, *Saccharomycodes ludwigii* or of the drink-spoiling molds of the genera *Mucor*, *Byssochlamys*, *Neosartorya*, *Aspergillus* und *Talaromyces*, in particular of the species *Mucor racemosus*, *Byssochlamys nivea*, *Neosartorya fischeri*, *Aspergillus fumigatus* und  
10 *A. fischeri*, *Talaromyces flavus*, *T. bacillisporus* and *T. flavus* or of the drink-spoiling bacteria of the genera *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Weissella*, *Lactococcus*, *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter*, *Bacillus* and *Alicyclobacillus*, in particular of the species *Lactobacillus collinoides*, *Leuconostoc mesenteroides*, *L. pseudomesenteroides*, *Oenococcus*  
15 *oeni*, *Bacillus coagulans*, *Alicyclobacillus ssp.*, *A. acidoterrestris*, *A. cycloheptanicus* and *A. herbarius*. In this context "specific hybridization" means that under the hybridization conditions described here or those known to the person skilled in the art in relation to *in situ* hybridization techniques, only the ribosomal RNA of the target organisms binds to the oligonucleotide, but not the  
20 rRNA of non-target microorganisms.

b) nucleic acid molecules which specifically hybridize under stringent conditions to a sequence complementary to the nucleic acid molecules mentioned in a) or to one of the probes SEQ ID No. 1, 5 to 146, 148 to 154, 157 to 160, 163 to 1033,  
25 1037 to 1138, 1142 to 1144.

c) Nucleic acid molecules comprising an oligonucleotide sequence of SEQ ID No. 1, 5 to 146, 148 to 154, 157 to 160, 163 to 1033, 1037 to 1138, 1142 to 1144 or the sequence of a nucleic acid molecule according to a) or b) and having at  
30 least one further nucleotide in addition to the mentioned sequences and their

derivatives, respectively, according to a) or b) and allowing specific hybridization with nucleic acid sequences of target organisms.

5 A further subject of the invention are also derivatives of the above competitor probe sequences, showing specific hybridizations with target nucleic acid sequences of the respective non-target genera and species, respectively, despite variations in sequence and/or length, and which therefore prevent the binding of the oligonucleotide probe to the nucleic acid sequences of the genera and species, respectively, not to be detected. They are suitable for use in a method according to  
10 the invention and ensure a specific detection of the respective microorganism. These derivatives especially include

a) nucleic acid molecules which (i) are identical in terms of bases to one of the above oligonucleotide sequences (SEQ ID No. 2 to 4, 147, 155 to 156, 161 to 162, 1034 to  
15 1036, 1139 to 1141) to at least 80%, preferably to at least 90%, particularly preferably to at least 92%, 94%, 96%, or (ii) differ from the above oligonucleotide sequences by one or more deletions and/or additions and which inhibit a specific hybridization of a specific oligonucleotide probe to nucleic acid sequences of a microorganism not to be detected.

20

b) Nucleic acid molecules which specifically hybridize to a sequence complementary to the nucleic acid molecules mentioned in a) or to one of the probes SEQ ID No. 2 to 4, 147, 155 to 156, 161 to 162, 1034 to 1036, 1139 to 1141 under stringent conditions.

25

c) Nucleic acid molecules comprising an oligonucleotide sequence of SEQ ID No. 2 to 4, 147, 155 to 156, 161 to 162, 1034 to 1036, 1139 to 1141 or the sequence of a nucleic acid molecule according to a) or b) and having at least one further nucleotide in addition to the mentioned sequences and their derivatives, respectively, according

to a) or b) and prevent the binding of a specific oligonucleotide probe to the nucleic acid sequence of a non-target microorganism.

5 The degree of sequence identity of a nucleic acid probe molecule to the oligonucleotide probes having SEQ ID No. 1 to SEQ ID No. 1144 can be determined using the usual algorithms. In this respect, for example, the program for determining the sequence identity available under <http://www.ncbi.nlm.nih.gov/BLAST> (on this page for example the link "Standard nucleotide-nucleotide BLAST [blastn]") is suitable.

10

In the present invention „hybridization“ can have the same meaning as „complementary“. The present invention also comprises those oligonucleotides, which hybridize to the (theoretical) antisense strand of one of the inventive oligonucleotides including the derivatives of the present invention of SEQ ID No. 1  
15 bis SEQ ID No. 1144.

The term “stringent conditions“ generally means conditions under which a nucleic acid sequence preferentially hybridizes to its target sequence and to a clearly lower extent, or not at all, to other sequences. Stringent conditions are partly sequence-  
20 dependent and will vary under different circumstances. Longer sequences hybridize specifically at higher temperatures. In general, stringent conditions are selected in such a way that the temperature is approximately 5°C below the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength, pH and nucleic acid concentration. The  $T_m$  is the temperature (under defined ionic strength, pH and  
25 nucleic acid concentration) at which 50% of the probe molecules complementary to the target sequence hybridize to the target sequence in the steady state.

The nucleic acid probe molecules of the present invention may be used within the detection method with various hybridization solutions. Various organic solvents may  
30 be used in concentrations of 0-80%. By keeping stringent hybridization conditions, it

is guaranteed that the nucleic acid probe molecule indeed hybridizes to the target sequence. Moderate conditions within the meaning of the invention are e.g. 0% formamide in a hybridization buffer as described below. Stringent conditions within the meaning of the invention are for example 20 % to 80 % formamide in the hybridization buffer.

Within the method according to the invention for the specific detection of yeasts of the genera *Zygosaccharomyces*, *Hanseniaspora*, *Candida*, *Brettanomyces*, *Dekkera*, *Pichia*, *Saccharomyces* and *Saccharomycodes*, in particular of the species *Zygosaccharomyces bailii*, *Z. mellis*, *Z. rouxii*, *Z. bisporus*, *Z. fermentati*, *Z. microellipsoides*, *Hanseniaspora uvarum*, *Candida intermedia*, *C. crusei* (*Issatchenkia orientalis*), *C. parapsilosis*, *Brettanomyces bruxellensis*, *B. naardenensis*, *Dekkera anomala*, *Pichia membranaefaciens*, *P. minuta*, *P. anomala*, *Saccharomyces exiguus*, *S. cerevisiae*, *Saccharomycodes ludwigii* a typical hybridization solution contains 0%-80% formamide, preferably 20%-60% formamide, particularly preferably 40% formamide. In addition, it has a salt concentration of 0.1 mol/l – 1.5 mol/l, preferably of 0.7 mol/l – 1.0 mol/l, and particularly preferably of 0.9 mol/l, whereby the salt preferably being sodium chloride. Further, the hybridization solution usually comprises a detergent, such as for instance sodium dodecyl sulfate (SDS) in a concentration of 0.001% - 0.2%, preferably in a concentration of 0.005% – 0.05%, particularly preferably in a concentration of 0.01%. For buffering the hybridization solution, various compounds such as Tris-HCl, sodium citrate, PIPES or HEPES may be used, which are usually used in concentrations of 0.01 – 0.1 mol/l, preferably of 0.01 to 0.05 mol/l, in a pH range of 6.0 – 9.0, preferably 7.0 to 8.0. The particularly preferred embodiment of the hybridization solution in accordance with the invention contains 0.02 mol/l Tris-HCl, pH 8.0.

Within the method according to the invention for the specific detection of molds of the genera *Mucor*, *Byssoschlamys*, *Neosartorya*, *Aspergillus* and *Talaromyces*, in

particular of the species *Mucor racemosus*, *Byssoschlamys nivea*, *Neosartorya fischeri*, *Aspergillus fumigatus* und *A. fischeri*, *Talaromyces flavus*, *T. bacillisporus* and *T. flavus*, a typical hybridization solution contains 0%-80% formamide, preferably 10%-60% formamide, particularly preferably 20% formamide. In addition, it has a salt concentration of 0.1 mol/l – 1.5 mol/l, preferably of 0.7 mol/l – 1.0 mol/l, and particularly preferably of 0.9 mol/l, whereby the salt preferably being sodium chloride. Further, the hybridization solution usually comprises a detergent, such as for instance sodium dodecyl sulfate (SDS) at a concentration of 0.001% - 0.2%, preferably at a concentration of 0.005 – 0.05%, particularly preferably at a concentration of 0.01%. For buffering the hybridization solution, various compounds such as Tris-HCl, sodium citrate, PIPES or HEPES may be used, which are usually used in concentrations of 0.01 – 0.1 mol/l, preferably of 0.01 to 0.05 mol/l, in a pH range of 6.0 – 9.0, preferably 7.0 to 8.0. The particularly preferred embodiment of the hybridization solution in accordance with the invention contains 0.02 mol/l Tris-HCl, pH 8.0.

Within the method according to the invention for the specific detection of bacteria of the genera *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Weissella*, *Lactococcus*, *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter*, *Bacillus* and *Alicyclobacillus*, in particular of the species *Lactobacillus collinoides*, *Leuconostoc mesenteroides*, *L. pseudomesenteroides*, *Oenococcus oeni*, *Bacillus coagulans*, *Alicyclobacillus ssp.*, *A. acidoterrestris*, *A. cycloheptanicus* and *A. herbarius*, a typical hybridization solution contains 0%-80% formamide, preferably 10%-60% formamide, particularly preferably 20% formamide. In addition, it has a salt concentration of 0.1 mol/l – 1.5 mol/l, preferably of 0.7 mol/l – 1.0 mol/l, and particularly preferably of 0.9 mol/l, whereby the salt preferably being sodium chloride. Further, the hybridization solution usually comprises a detergent, such as for instance sodium dodecyl sulfate (SDS) at a concentration of 0.001% - 0.2%, preferably at a concentration of 0.005% – 0.05%, particularly preferably at a concentration of 0.01%. For buffering the hybridization solution, various compounds such as Tris-HCl, sodium citrate, PIPES



or HEPES may be used, which are usually used in concentrations of 0.01 – 0.1 mol/l, preferably of 0.01 to 0.05 mol/l, in a pH range of 6.0 – 9.0, preferably 7.0 to 8.0. The particularly preferred embodiment of the hybridization solution in accordance with the invention contains 0.02 mol/l Tris-HCl, pH 8.0.

5

It shall be understood that the one skilled in the art can select the specified concentrations of the constituents of the hybridization buffer in such a way that the desired stringency of the hybridization reaction is achieved. Particularly preferred embodiments are related from stringent to particularly stringent hybridization conditions. Using these stringent conditions the one skilled in the art can determine whether a particular nucleic acid molecule allows the specific detection of nucleic acid sequences of target organisms and may thus be reliably used within the invention.

15 The concentration of the nucleic acid probe in the hybridization buffer depends on the kind of label and on the number of target structures. In order to allow rapid and efficient hybridization, the number of nucleic acid probe molecules should exceed the number of target structures by several orders of magnitude. However, it has to be taken into consideration that in fluorescence *in situ*-hybridization (FISH) too high levels of fluorescently labelled nucleic acid probe molecules result in increased background fluorescence. The concentration of the nucleic acid probe molecules should therefore be in the range between 0.5 and 500 ng/μl. Within the method of the present invention the preferred nucleic acid probe concentration is between 1.0 and 10 ng for each nucleic acid probe molecule used per μl of hybridization solution. The volume of hybridization solution used should be between 8 μl and 100 ml, in a particularly preferred embodiment of the method of present invention it is 30 μl.

The concentration of the competitor probe in the hybridization buffer depends on the number of target structures. In order to allow rapid and efficient hybridization, the number of competitor probes should exceed the number of target structures by

30

several orders of magnitude. The concentration of the competitor probe molecules should therefore be in a range between 0.5 and 500 ng/ $\mu$ l. Within the method of the present invention the preferred concentration is between 1.0 and 10 ng for each competitor probe molecule used per  $\mu$ l of hybridization solution. The volume of hybridization solution used should be between 8  $\mu$ l and 100 ml, in a particularly preferred embodiment of the method of present invention it is 30  $\mu$ l.

The hybridization usually lasts between 10 minutes and 12 hours, preferably the hybridization lasts for about 1.5 hours. The hybridization temperature is preferably between 44°C and 48°C, particularly preferably 46°C, whereby the parameter of the hybridization temperature as well as the concentration of salts and detergents in the hybridization solution may be optimized depending on the nucleic acid probes, especially their lengths and the degree to which they are complementary to the target sequence in the cell to be detected. The one skilled in the art is familiar with appropriate calculations.

After hybridization the non-hybridized and excess nucleic acid probe molecules should be removed or washed off, which is usually achieved by a conventional washing solution. This washing solution may, if desired, contain 0.001-0.1%, preferably 0.005-0.05%, particularly preferably 0.01% of a detergent such as SDS, as well as Tris-HCl in a concentration of 0.001-0.1 mol/l, preferably 0.01-0.05 mol/l, particularly preferably 0.02 mol/l, wherein the pH value of Tris-HCl is within the range of 6.0 to 9.0, preferably of 7.0 to 8.0, particularly preferably 8.0. A detergent may be contained, although this is not obligatorily necessary. Furthermore, the washing solution usually contains NaCl, whereby the concentration is 0.003 mol/l to 0.9 mol/l, preferably 0.01 mol/l to 0.9 mol/l, depending on the stringency required. Moreover, the washing solution may contain EDTA, whereby the concentration is preferably 0.005 mol/l. The washing solution may further contain suitable amounts of preservatives known to the expert.

In general, buffer solutions are used in the washing step which can in principle be very similar to the hybridization buffer (buffered sodium chloride solution), except that the washing step is usually performed in a buffer with a lower salt concentration and at a higher temperature, respectively. For theoretical estimation of the hybridization conditions, the following formula may be used:

$$T_d = 81.5 + 16.6 \lg[\text{Na}^+] + 0.4 \times (\% \text{ GC}) - 820/n - 0.5 \times (\% \text{ FA})$$

$T_d$  = dissociation temperature in °C

10  $[\text{Na}^+]$  = molarity of the sodium ions

% GC = percentage of guanine and cytosine nucleotides relative to the total number of bases

$n$  = length of the hybrid

% FA = formamide content

15

Using this formula, the formamide content (which should be as low as possible due to the toxicity of the formamide) of the washing buffer may for example be replaced by a correspondingly lower sodium chloride content. However, the person skilled in the art is, from the extensive literature concerning in situ hybridization methods, aware of the fact that, and in which way, the mentioned contents can be varied.

20

Concerning the stringency of the hybridization conditions, the same applies as outlined above for the hybridization buffer.

25

The “washing off” of the non-bound nucleic acid probe molecules is usually performed at a temperature in the range of 44°C to 52°C, preferably of 44°C to 50°C and particularly preferably at 46°C for 10 to 40 minutes, preferably for 15 minutes.

30

The specifically hybridized nucleic acid probe molecules can then be detected in the respective cells, provided that the nucleic acid probe molecule is detectable, e.g., by linking the nucleic acid probe molecule to a marker by covalent binding. As

detectable markers, for example, fluorescent groups, such as for example CY2 (available from Amersham Life Sciences, Inc., Arlington Heights, USA), CY3 (also available from Amersham Life Sciences), CY5 (also obtainable from Amersham Life Sciences), FITC (Molecular Probes Inc., Eugene, USA), FLUOS (available from Roche Diagnostics GmbH, Mannheim, Germany), TRITC (available from Molecular Probes Inc., Eugene, USA), 6-FAM or FLUOS-PRIME are used, which are well known to the person skilled in the art. Also chemical markers, radioactive markers or enzymatic markers, such as horseradish peroxidase, acid phosphatase, alkaline phosphatase, peroxidase may be used. For each of these enzymes a number of chromogens is known which may be converted instead of the natural substrate and may be transformed into either coloured or fluorescent products. Examples of such chromogens are listed in the following table:

**Table**

15

Enzyme	Chromogen
1. Alkaline phosphatase and acid phosphatase	4-methylumbelliferyl phosphate (*), bis(4-methylumbelliferyl phosphate, (*) 3-O-methylfluorescein, flavone-3-diphosphate triammonium salt (*), p-nitrophenylphosphate disodium salt
2. Peroxidase	tyramine hydrochloride (*), 3-(p-hydroxyphenyl)-propionate (*), p-hydroxyphenethyl alcohol (*), 2,2'-azino-di-3-ethylbenzothiazoline sulfonic acid (ABTS), ortho-phenylendiamine dihydrochloride, o-dianisidine, 5-aminosalicylic acid, p-ucresol (*), 3,3'-dimethyloxy benzidine, 3-methyl-2-benzothiazoline hydrazone, tetramethylbenzidine

- |                           |  |
|---------------------------|--|
| 3. Horseradish peroxidase | H <sub>2</sub> O <sub>2</sub> + diammonium benzidine<br>H <sub>2</sub> O <sub>2</sub> + tetramethylbenzidine |
| 4. β-D-galactosidase      | o-nitrophenyl-β-D-galactopyranoside, 4-methylumbelliferyl-β-D-galactoside                                    |
| 5. Glucose oxidase        | ABTS, glucose and thiazolyl blue   |

\* fluorescence

5 Finally, it is possible to design the nucleic acid probe molecules in such a way that another nucleic acid sequence suitable for hybridization is present at their 5' or 3' ends. This nucleic acid sequence in turn comprises about 15 to 100, preferably 15-50 nucleotides. This second nucleic acid region may in turn be detected by a nucleic acid probe molecule which is detectable by one of the above-mentioned agents.

10 Another possibility is the coupling of the detectable nucleic acid probe molecules to a haptene which may subsequently be brought into contact with an antibody recognising the haptene. Digoxigenin may be mentioned as an example of such a haptene. Other examples in addition to those mentioned are well known to the one skilled in the art.

15

The final evaluation is, depending on the kind of labelling of the probe used, possible, among others, with an optical microscope, epifluorescence microscope, chemoluminometer, fluorometer.

20 An important advantage of the methods described in this application for the specific detection of drink-spoiling yeasts of the genera *Zygosaccharomyces*, *Hanseniaspora*, *Candida*, *Brettanomyces*, *Dekkera*, *Pichia*, *Saccharomyces* and *Saccharomycodes*, in particular of the species *Zygosaccharomyces bailii*, *Z. mellis*, *Z. rouxii*, *Z. bisporus*, *Z. fermentati*, *Z. microellipsoides*, *Hanseniaspora uvarum*, *Candida intermedia*, *C. crusei* (*Issatchenkia orientalis*), *C. parapsilosis*, *Brettanomyces bruxellensis*, *B.*

25

*naardenensis*, *Dekkera anomala*, *Pichia membranaefaciens*, *P. minuta*, *P. anomala*, *Saccharomyces exiguus*, *S. cerevisiae*, *Saccharomycodes ludwigii* or for the specific detection of drink-spoiling molds of the genera *Mucor*, *Byssoschlamys*, *Neosartorya*, *Aspergillus* and *Talaromyces*, in particular of species *Mucor racemosus*,  
5 *Byssoschlamys nivea*, *Neosartorya fischeri*, *Aspergillus fumigatus* and *A. fischeri*, *Talaromyces flavus*, *T. bacillisporus* and *T. flavus*, or for the specific detection of drink-spoiling bacteria of the genera *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Weissella*, *Lactococcus*, *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter*, *Bacillus* und *Alicyclobacillus*, in particular of the species *Lactobacillus collinoides*,  
10 *Leuconostoc mesenteroides*, *L. pseudomesenteroides*, *Oenococcus oeni*, *Bacillus coagulans*, *Alicyclobacillus ssp.*, *A. acidoterrestris*, *A. cycloheptanicus* and *A. herbarius* compared to the detection methods described above is the exceptional speed. In comparison to conventional cultivation methods which need up to 10 days, the result is obtained within 24 to 48 hours when the methods according to the  
15 invention are used.

Another advantage is the ability to perform an accurate differentiation of the drink-spoiling microorganisms to be detected. With the methods common up to now no differentiation of the microorganisms was carried out until the genus or species level,  
20 as the differentiation was either not possible at all or was too time-consuming.

Another advantage is the specificity of these methods. With the nucleic acid probe molecules used, drink-spoiling yeasts of the genera *Zygosaccharomyces*, *Hanseniaspora*, *Candida*, *Brettanomyces*, *Dekkera*, *Pichia*, *Saccharomyces* and  
25 *Saccharomycodes*, in particular the species *Zygosaccharomyces bailii*, *Z. mellis*, *Z. rouxii*, *Z. bisporus*, *Z. fermentati*, *Z. microellipsoides*, *Hanseniaspora uvarum*, *Candida intermedia*, *C. crusei* (*Issatchenkia orientalis*), *C. parapsilosis*, *Brettanomyces bruxellensis*, *B. naardensis*, *Dekkera anomala*, *Pichia membranaefaciens*, *P. minuta*, *P. anomala*, *Saccharomyces exiguus*, *S. cerevisiae*,  
30 *Saccharomycodes ludwigii* or drink-spoiling molds of the genera *Mucor*,

*Byssochlamys*, *Neosartorya*, *Aspergillus* and *Talaromyces*, in particular of the species *Mucor racemosus*, *Byssochlamys nivea*, *Neosartorya fischeri*, *Aspergillus fumigatus* and *A. fischeri*, *Talaromyces flavus*, *T. bacillisporus* and *T. flavus* or drink-spoiling bacteria of the genera *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Weissella*,  
5 *Lactococcus*, *Acetobacter*, *Gluconobacter*, *Gluconoacetobacter*, *Bacillus* and *Alicyclobacillus*, in particular of the species *Lactobacillus collinoides*, *Leuconostoc mesenteroides*, *L. pseudomesenteroides*, *Oenococcus oeni*, *Bacillus coagulans*, *Alicyclobacillus ssp.*, *A. acidoterrestris*, *A. cycloheptanicus* and *A. herbarius* can be detected in a highly specific manner. By the visualisation of the microorganisms a  
10 visual control may be performed at the same time. False-positive results, such as often occurring in polymerase chain reaction, are therefore ruled out.

Another advantage of the methods according to the invention is their ease of use. Thus, using this methods, large numbers of samples can be easily tested regarding  
15 the presence of the mentioned microorganisms.

Finally, an important advantage compared to the state of the art is the possible simultaneous detection of several of the mentioned microorganisms by the use of respective mixtures of probes. Following this approach all practise relevant drink-spoiling microorganisms can be detected in a few tests.  
20

Different probes may hereby be coupled with different labels, so that the various, detected micororganisms may be discriminated in an easy and reliable way. For example, a first oligonucleotide may be specifically labelled with a green  
25 fluorescence dye and serves for the detection of a certain genus or species of microorganism. A second oligonucleotide is also specifically labelled with, for instance, a red fluorescence dye and serves for the detection of a second genus or species of microorganism. The oligonucleotides referred to as competitor probes remain non-labelled and prevent the binding of the first and/or the second  
30 oligonucleotide probe to bacteria which do not belong to the genera or species to be

detected. The different labels, e.g. the green fluorescence dye on the one hand and the red fluorescence dye on the other hand may be differentiated in an easy manner, for example by using different filters in fluorescence microscopy.

- 5 The methods according to the invention may be used in various ways.

For example, non-alcoholic drinks (e.g. fruit juices, fruit nectars, fruit concentrates, mashed fruits, soft drinks and waters) may be tested for the presence of the microorganisms to be detected.

10

For example, also environmental samples can be tested for the presence of the micororganisms to be detected. Theses samples may be, for example, collected from soil or be parts of plants.

- 15 The method according to the invention may further be used for testing sewage samples or silage samples.

The method according to the invention may further be used for testing medicinal samples, e.g. stool samples, blood cultures, sputum, tissue samples (also sections),  
20 wound material, urine, samples from the respiratory tract, implants and catheter surfaces.

Another field of use of the method according to the invention is the control of food. In preferred embodiments the food samples are obtained from milk or milk products  
25 (yogurt, cheese, curd, butter, buttermilk), drinking water, alcoholic drinks (beer, wine, spirits), bakery products or meat products.

A further field of use of the method according to the invention is the analysis of pharmaceutical and cosmetic products, e.g. ointments, creams, tinctures, juices,  
30 solutions, drops, etc.



Furthermore, according to the invention, kits for performing the respective methods are provided. The hybridization arrangement contained in these kits is described for example in German patent application 100 61 655.0. Express reference is herewith  
5 made to the disclosure contained in this document with respect to the in situ hybridization arrangement.

Besides the described hybridization arrangement (referred to as VIT reactor), the most important component of the kits is the respective hybridization solution  
10 (referred to as VIT solution) with the nucleic acid probe molecules specific for the microorganisms to be detected, which are described above (VIT solution). Further contained are the respective hybridization buffer (Solution C) and a concentrate of the respective washing solution (Solution D). Also contained are optionally fixation  
15 solutions (Solution A and Solution B) as well as optionally an embedding solution (finisher). Optionally, solutions are contained for performing in parallel a positive control as well as of a negative control.

The following example is intended to illustrate the invention without limitation.

## 20 Example

### Specific rapid detection of drink-spoiling microorganisms in a sample

A sample is cultivated for 20 to 48 hours in a suitable manner. For the detection of  
25 yeasts and molds cultivation may be performed, for example, in SSL-bouillon for 24 hours at 25°C. For the detection of lactic acid bacteria the cultivation may be performed for example in MRS-bouillon for 48 hours at 30°C. For the detection of acetate acid bacteria the cultivation may be performed, for example, on DSM-agar for 48 hours at 28°C. For the detection of bacilli, in particular *B. coagulans*, the  
30 cultivation may be performed, for example, on dextrose-casein-peptone-agar for 48

hours at 55°C. For the detection of alicyclobacilli, the cultivation may be performed, for example, in BAM-bouillon for 48 hours at 44°C.

5 To an aliquot of the culture the same volume of fixation solution (Solution B, ethanol absolute) is added. Alternatively, an aliquot of the culture may be centrifuged (4000 g, 5 min, room temperature) and, after discarding the supernatant, the pellet may be dissolved in 4 drops of fixation solution (Solution B).

10 For performing the hybridization a suitable aliquot of the fixed cells (preferably 5 µl) is applied onto a slide and dried (46°C, 30 min, or until completely dry). Alternatively, the cells may also be applied to other carrier materials (e.g. a microtiter plate or a filter). The dried cells are then completely dehydrated by again adding the fixation solution (Solution B). The slide is again dried (room temperature, 3 min, or until completely dry).

15 Then the hybridization solution (VIT solution, hybridization buffer containing labeled probe molecules) containing the above described nucleic acid probe molecules specific for the microorganisms to be detected, is applied to the fixed, dehydrated cells. The preferred volume is 40 µl. The slide is then incubated (46°C, 20 90 min) in a chamber humidified with hybridization buffer (Solution C), preferably the VIT reactor (c. f. DE 100 61 655.0).

25 Then the slide is removed from the chamber, the chamber is filled with washing solution (Solution D, diluted 1:10 with distilled water) and the slide is incubated in the chamber (46°C, 15 min).

Then the chamber is filled with distilled water, the slide is briefly immersed and then air-dried in lateral position (46°C, 30 min or until completely dry).

30 Then the slide is embedded in a suitable medium (Finisher).

Finally, the sample is analyzed with the help of a fluorescence microscope.